

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number:

0 412 404 A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90114628.2

(51) Int. Cl.⁵: **C07D 277/38**, A61K 31/425,
C07D 277/56, C07D 417/12

(22) Date of filing: 31.07.90

(30) Priority: 07.08.89 GB 8918045
21.02.90 GB 9003930

(43) Date of publication of application:
13.02.91 Bulletin 91/07

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: FUJISAWA PHARMACEUTICAL CO.,
LTD.
4-7, Doshomachi 3-chome Chuo-ku
Osaka-shi Osaka 541(JP)

(72) Inventor: Matsuo, Masaaki
4-12, Nakasakurazuka 5-chome
Toyonaka-shi, Osaka 560(JP)

Inventor: Ogino, Takashi
4-6-208, Minamigoyo 2-chome
Kita-ku, Kobe-shi, Hyogo 651-11(JP)


Inventor: Igari, Norihiro
2-12-401 Ohsho-kita 4-chome
Amagasaki-shi, Hyogo 660(JP)

Inventor: Seno, Hachiro
12-1, Sengokuhigashimachi
Kadoma-shi, Osaka 571(JP)
Inventor: Shimomura, Kyoichi
3-9, Tsugumodai 2-chome
Suita-shi, Osaka 565(JP)

(74) Representative: Türk, Gille, Hrabal
Brucknerstrasse 20
D-4000 Düsseldorf 13(DE)

(54) Thiazole derivatives, processes for production thereof and pharmaceutical compositions comprising the same.

(57) A compound of the formula :




wherein

R¹ is hydrogen or acyl which may be substituted with halogen,

R² is hydrogen, lower alkyl, hydroxy(lower)alkyl, halogen or carboxy,

A is -CH₂-, -CO-, -C(=NOR⁴)- [wherein R⁴ is lower alkyl,



[wherein m is 0, 1 or 2], and

R³ is aryl which may be substituted with halogen, hydroxy, lower alkoxy, nitro, amino or acylamino; or
N-containing unsaturated heterocyclic group which may be substituted with lower alkyl, amino, hydroxy or halo-

EP 0 412 404 A2


(lower)alkyl,
and pharmaceutically acceptable salts thereof,
processes for their preparation and pharmaceutical compositions comprising them as an active ingredient in
admixture with pharmaceutically acceptable carriers.

More particularly, this invention relates to new thiazole derivatives and pharmaceutically acceptable salts thereof which have pharmacological activities, processes for production thereof, pharmaceutical compositions comprising the same and methods of use thereof.

Another object of this invention is to provide processes for production of said thiazole derivatives and pharmaceutically acceptable salts thereof.

Still further object of this invention is to provide methods of using said thiazole derivatives or pharmaceutically acceptable salts thereof for therapeutic treatment of rheumatism, nephritis, thrombocytopenia, tumor or side effect of an antitumor agent in human being and animals.

The object thiazole derivatives of this invention are novel and represented by the following general
15 formula (I):




R¹ is hydrogen or acyl which may be substituted with halogen,
R² is hydrogen, lower alkyl, hydroxyl(lower)alkyl, halogen or carboxy,
A is -CH₂-, -CO-, -C(=NOR⁴)- [wherein R⁴ is lower alkyl],











R³ is aryl which may be substituted with halogen, hydroxy, lower alkoxy, nitro, amino or acylamino; or N-containing unsaturated heterocyclic group which may be substituted with lower alkyl, amino, hydroxy or halo(lower)alkyl.



40 The object compound (I) of the present invention can be prepared by the following processes.

Process 1



Process 2Process 3Process 4Process 5

Process 6Process 7Process 8Process 9

Process 10Process 11

wherein

R¹, R², R³, R⁴ and A are each as defined above,

R⁵ is acyl which may be substituted with halogen,

R⁶ is acylamino,

X is halogen,


P is 0 or 1,

n is 0 or 1 and

q is 1 or 2,

provided that q is 2 when n is 1.

A kind of the starting compounds [II] can be prepared by the process as illustrated in the following.

Process A

wherein

R¹ and x are each as defined above, and

R⁷ is lower alkyl or hydroxy(lower)alkyl.

55 In the above and subsequent description of the present specification, suitable examples and illustrations for the various definitions to be included within the scope of the invention are explained in detail as follows:

The term "lower" is intended to mean 1 to 6 carbon atom(s), unless otherwise indicated.

Suitable "lower alkyl" may be a straight or branched one such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl or the like.

Suitable example of "lower alkyl" moiety in the term "hydroxy(lower)alkyl" and "halo(lower)alkyl" can be referred to the ones as exemplified above.

Suitable "halo(lower)alkyl" may include "monohalo(lower)alkyl" [e.g. chloromethyl, bromomethyl, fluoromethyl, etc.], "dihalo(lower)alkyl" [e.g. dichloromethyl, dibromomethyl, difluoromethyl, etc.] and "trihalo(lower)alkyl" [e.g. trichloromethyl, tribromomethyl, trifluoromethyl, trifluoroethyl, etc.] and the like.

Suitable "lower alkoxy" may include methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, t-butoxy, pentyloxy, hexyloxy and the like.

Suitable "halogen" may include fluorine, chlorine, bromine and iodine.

Suitable examples of "aryl" may include phenyl, tolyl, xylyl, cumenyl, naphthyl, and the like.

Suitable acyl may include an aliphatic acyl, an aromatic acyl and an aliphatic acyl substituted with aromatic group(s).

The aliphatic acyl may include saturated or unsaturated, acyclic or cyclic ones, such as lower alkanoyl (e.g. formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, etc.), lower alkoxycarbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, tert-butoxycarbonyl, etc.), lower alkanesulfonyl [e.g. methanesulfonyl, ethanesulfonyl, propanesulfonyl, butanesulfonyl, pentanesulfonyl, hexanesulfonyl, etc.], lower alkenoyl (e.g. acryloyl, methacryloyl, crotonoyl, etc.), carbamoyl and the like.

The aromatic acyl may include aroyl (e.g. benzoyl, toluoyl, xyloyl, etc.) and the like.

The aliphatic acyl substituted with aromatic group(s) may include ar(lower)alkanoyl such as phenyl-(lower)alkanoyl (e.g. phenylacetyl, phenylpropionyl, phenylhexanoyl, etc.), ar(lower)alkoxycarbonyl such as phenyl(lower)alkoxycarbonyl (e.g. benzyloxycarbonyl, phenethylloxycarbonyl, etc.), phenoxy(lower)alkanoyl (e.g. phenoxyacetyl, phenoxypropionyl, etc.), and the like.

Suitable example of "acyl" moiety in the term of "acylamino" can be referred to the ones as exemplified above.

Suitable "N-containing unsaturated heterocyclic group" may be one containing at least one nitrogen atom and may include monocyclic or polycyclic heterocyclic group, and preferable heterocyclic group may be unsaturated 3 to 6 membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazolyl (e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.), tetrazolyl (e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.), etc.; unsaturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indolizyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl (e.g., tetrazolo-[1,5-b]pyridazinyl, etc.), etc.; unsaturated 3- to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.), etc.; unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g., benzoxazolyl, benzoxadiazolyl, etc.); unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl (e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.), etc.; unsaturated condensed heterocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., benzothiazolyl, benzothiadiazolyl, etc.) and the like. Said "heterocyclic group" may have 1 to 4 substituents such as lower alkyl as exemplified above.

Suitable pharmaceutically acceptable salts of the object compounds [I] are conventional non-toxic salts and include an organic acid salt [e.g. formate, acetate, trifluoroacetate, maleate, tartrate, methanesulfonate, benzenesulfonate, toluenesulfonate, etc.], an inorganic acid salt [e.g. hydrochloride, hydrobromide, sulfate, phosphate, etc.], a salt with an amino acid [e.g. arginine, glutamic acid, ornithine, etc.], a metal salt such as an alkali metal salt [e.g. sodium salt, potassium salt, etc.] and an alkaline earth metal salt [e.g. calcium salt, magnesium salt, etc.], an ammonium salt, an organic base salt [e.g. trimethylamine salt, triethylamine salt, pyridine salt, picoline salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt, etc.], and the like.

In this respect, it is to be noted that the compounds [Ia] to [Ip] are included within the scope of the compound [I], and accordingly the suitable salts of these compounds [Ia] to [Ip] are to be referred to those as exemplified for the object compounds [I] in the above.

The processes for preparing the object compound [I] or salts thereof are explained in detail in the following.

55 Process 1

The object compound [Ib] or its salt can be prepared by deacylating a compound [Ia] or its salt.

Suitable method for this deacylation reaction may include conventional one such as hydrolysis and the

like.

Hydrolysis is preferably carried out in the presence of an acid.

Suitable acid may be an inorganic acid [e.g. hydrochloric acid, hydrobromic acid, sulfuric acid, etc.], an organic acid [e.g. formic acid, acetic acid, trifluoroacetic acid, propionic acid, benzenesulfonic acid, p-toluenesulfonic acid, etc.], an acidic ion-exchange resin and the like. In case that the organic acid such as trifluoroacetic acid and p-toluenesulfonic acid is used in this reaction, the reaction is preferably carried out in the presence of cation trapping agents [e.g. anisole, etc.].

The acid suitable for this hydrolysis can be selected according to the kinds of the acyl group to be removed.

The hydrolysis is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, methanol, ethanol, propanol, tert-butyl alcohol, tetrahydrofuran, N,N-dimethylformamide, dioxane or a mixture thereof, and further the above-mentioned acids can also be used as a solvent when they are in liquid.

The reaction temperature of this hydrolysis is not critical, and the reaction is usually carried out under cooling, at ambient temperature or under heating.

In this process, when the starting compound [Ia] or its salt has a acylamino group for R³, the group is also converted to an amino group.

20 Process 2

The object compound [Ic] or its salt can be prepared by reacting a compound [II] or its salt with a compound [III] or its salt.

Suitable salts of the compound [III] and [III] may be the same as those exemplified as base salts of the object compound [I].

This reaction is usually carried out in a solvent such as methanol, ethanol, propanol, tetrahydrofuran, dioxane, dimethylformamide or any other organic solvent which does not adversely influence the reaction.

In case that a free form of the compound [III] is used in this reaction, the reaction is preferably carried out in the presence of a conventional base, such as alkali metal hydride [e.g. sodium hydride, potassium hydride, etc.] alkaline earth metal hydride [e.g. calcium hydride, magnesium hydride, etc.], alkali metal hydroxide [e.g. sodium hydroxide, potassium hydroxide, etc.], alkali metal carbonate [e.g. sodium carbonate, potassium carbonate, etc.], alkali metal bicarbonate [e.g. sodium bicarbonate, potassium bicarbonate, etc.], alkali metal fluoride [e.g. potassium fluoride, cesium fluoride, etc.], alkali metal alkoxide [e.g. sodium methoxide, sodium ethoxide, potassium tert-butoxide, etc.], trialkylamine [e.g. trimethylamine, triethylamine, etc.], picoline, 1,5-diazabicyclo[4,3,0]non-5-ene, 1,4-diazabicyclo[2,2,2]-octane, 1,5-diazabicyclo[5,4,0]undecene-5 or the like.

The reaction temperature is not critical, and the reaction is usually carried out at ambient temperature or under cooling, warming or heating.

40 Process 3

The object compound [Ie] or its salt can be prepared by reducing a compound [Id] or its salt.

The reduction can be carried out in a conventional manner, namely, chemical reduction or catalytic reduction.

Suitable reducing agents to be used in chemical reduction are a combination of metal (e.g. stannum, zinc, iron, etc.) and ammonium chloride or an base (e.g. ammonia, sodium hydroxide, etc.), a combination of metal (e.g. tin, zinc, iron, etc.) or metallic compound (e.g. chromium chloride, stannous chloride, chromium acetate, etc.) and an organic or inorganic acid (e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.), alkali metal borohydride (e.g. lithium borohydride, sodium borohydride, potassium borohydride, etc.) alkali metal cyanoborohydride (e.g. sodium cyanoborohydride, etc.) or alkali metal ammonium hydride (e.g. lithium aluminum hydride, etc.) or the like.

Suitable catalysts to be used in catalytic reduction are conventional ones such as platinum catalyst (e.g. platinum plate, spongy platinum, platinum black, colloidal platinum, platinum oxide, platinum wire, etc.), palladium catalyst (e.g. palladium on carbon, colloidal palladium, palladium on barium sulfate, palladium on barium carbonate, etc.), nickel catalyst (e.g. reduced nickel, nickel oxide, Raney nickel, etc.), cobalt catalyst (e.g. reduced cobalt, Raney cobalt, etc.), iron catalyst (e.g. reduced iron, Raney iron, etc.), copper catalyst

(e.g. reduced copper, Raney copper, Ullman copper, etc.) or the like.

The reduction of this process is usually carried out in a solvent such as water, alcohol (e.g. methanol, ethanol, propanol, etc.), acetic acid, dioxane, tetrahydrofuran, N,N-dimethylformamide, dimethyl sulfoxide, or any other organic solvent which does not adversely influence the reaction, or a mixture thereof. In case

The reaction is preferably carried out under cooling to warming or heating.

Process 4

A compound [lg] or its salt can be prepared by subjecting a compound [lf] or its salt to oxidation.

Oxidation in this process is carried out in a conventional manner with a conventional oxidizing agent which can oxidize a -S- group into -SO- or -SO₂- group, or -SO- group into -SO₂- group.

Suitable example of such oxidizing agent are inorganic peracid or its salt (e.g. periodic acid, persulfuric acid, etc.) or the sodium or potassium salt thereof, an organic peracid or its salt (e.g. perbenzoic acid, 3-chloroperbenzoic acid, performic acid, peracetic acid, chloroperacetic acid, trifluoroperacetic acid, etc. or the sodium or potassium salt thereof, etc.), ozone, hydrogen peroxide, urea-hydrogen peroxide and the like.

The present reaction is preferably conducted in the presence of a compound comprising a Group Vb or Vlb metal in the Periodic Table, for example, tungstic acid, molybdic acid, vanadic acid, etc. or their salt with an alkali metal (e.g. sodium, potassium, etc.), an alkaline earth metal (e.g. calcium, magnesium, etc.) or ammonium, etc., or vanadium pentoxide.

The present oxidation is usually conducted in a solvent such as water, acetic acid, ethyl acetate, chloroform, dichloromethane, tetrahydrofuran, dioxane, N,N-dimethylformamide or any other solvent which does not give bad influence to the present reaction.

There is not particular limitation to the reaction temperature, and the present reaction is usually conducted at ambient temperature or under cooling.

Process 5

The object compound [li] or its salt can be prepared by subjecting a compound [lh] or its salt to oxidation.

Suitable oxidizing agent may include hypochlorite compound (e.g. tert-butyl hypochlorite, etc.), permanganate (e.g. potassium permanganate, etc.), or any other conventional oxidizing agent.

This reaction is usually carried out in a solvent such as water, acetic acid, diethyl ether, dioxane, dichloromethane, chloroform, N,N-dimethylformamide, dimethyl sulfoxide, or any other organic solvent which does not adversely influence the reaction, or a mixture thereof. In case the above-mentioned acid is liquid, it can be also used as a solvent.

The reaction can be carried out under cooling to warming or heating.

Process 6

The object compound [lk] or its salt can be prepared by halogenating a compound [lj] or its salt.

Suitable halogenating agent of this reaction may include conventional ones for example, halogen [e.g. chlorine, bromine, iodine, etc.], sulfuryl halide [e.g. sulfuryl chloride, sulfuryl bromide, etc.], N-halosuccinimide [e.g. N-chlorosuccinimide, N-bromosuccinimide, etc.], pyridinium hydrohalide perhalide [e.g. pyridinium hydrobromide perbromide, pyridinium hydrochloride perchloride, etc.], quaternary ammonium perhalide [e.g. phenyltrimethylammonium perbromide, etc.], ω-trihaloacetophenone [e.g. ω-tribromoacetophenone, etc.], cupric or potassium bromide, selenium oxychloride, or the like. These halogenating agents may be selected according to the kind of the starting compound [lj] to be used.

This reaction is usually carried out in a conventional solvent such as chloroform, methylene chloride, carbon tetrachloride, acetic acid, a mixture of hydrogen halide [e.g. hydrogen bromide, hydrogen chloride, etc.] and acetic acid, water, dimethylformamide or the like.

The reaction temperature is not critical, and the reaction is usually carried out under cooling, at ambient temperature or under warming or heating.

Process 7

The object compound [Ia] or its salt can be prepared by acylating a compound [Ib] or its reactive derivatives at the amino group or a salt thereof.

5 Suitable reactive derivatives at the amino group of the compound. [Ib] include conventional ones used in amidation for example, Schiff's base type imino or its tautomeric enamine type isomer formed by reaction of a compound [Ib] with a carbonyl compound, a silyl derivative formed by reaction of a compound [Ib] with a silyl compound such as trimethylsilylacetamide, bis(trimethylsilyl)acetamide or the like, a derivative formed by reaction of a compound [Ib] with phosphorus trichloride or phosgene, and the like.

10 Suitable acylating agent to be used in this reaction includes an organic acid such as alkanolic acid [e.g. formic acid, acetic acid, propionic acid, etc.], arenecarboxylic acid (e.g. benzoic acid, toluenecarboxylic acid, etc.) which may have halogen, lower alkanesulfonic acid [e.g. methanesulfonic acid, etc.], arylisocyanate [e.g. phenylisocyanate, etc.] which may have halogen and their reactive derivative.

The suitable reactive derivative may be a conventional one such as an acid halide [e.g. acid chloride, 15 acid bromide, etc.], an acid azide, an acid anhydride, an activated amide, an activated ester and the like. When free acid is used as an acylating agent, the acylation reaction may preferably be conducted in the presence of a conventional condensing agent such as N,N'-dicyclohexylcarbodiimide or the like.

This reaction is usually carried out in a solvent which does not adversely influence the reaction such as water, tetrahydrofuran, chloroform, dioxane, pyridine, methylene chloride, N,N-dimethylformamide or the 20 like.

The reaction temperature is not critical and the reaction can be carried out at any temperature under cooling to heating.

25 Process 8

The object compound [II] or its salt can be prepared by acylating a compound [Ie] or its reactive derivatives at the amino group or a salt thereof.

Suitable reactive derivatives at the amino group are to be referred to those as exemplified in Process 7.

30 This reaction may be carried out in substantially the same manner as Process 7, and therefore the reaction mode and reaction conditions [e.g. acylating agent, solvent, reaction temperature, etc.] of this reaction are to be referred to those as explained in Process 7.

35 Process 9

The object compound [Im] or its salt can be prepared by reacting a compound [IV] or its salt with a compound [V] or its salt.

40 Suitable salts of the compounds [IV] and [V] may be the same as those exemplified for the compound [I].

Suitable examples of the compound [V] may be thiocarbonyl derivatives such as thiourea, N-acylthiourea [e.g. N-formylthiourea, N-acetylthiourea, N-propionylthiourea, N-benzoylthiourea which may be substituted with halogen, etc.] or the like.

This reaction is usually carried out in a conventional solvent such as water, methanol, ethanol, isopropyl 45 alcohol, tetrahydrofuran, dioxane, chloroform, methylene chloride, dimethylacetamide, dimethylformamide or any other organic solvent which does not adversely influence the reaction. Among these solvents, hydrophilic solvents may be used in a mixture with water.

The reaction temperature is not critical, and the reaction is usually carried out at ambient temperature or under warming or heating.

50

Process 10

55 The object compound [In] or its salt can be prepared by reacting a compound [VI] or its salt with a compound [V] or its salt.

Suitable salts of the compound [VI] may be the same as those exemplified for the object compound [I].

This reaction may be carried out in substantially the same manner as Process 9, and therefore the reaction mode and reaction conditions [e.g. solvent, reaction temperature, etc.] of this reaction are to be

referred to those as explained in Process 9.

Process 11

5

The object compound [Ip] or its salt can be prepared by reacting a compound [Io] or its salt with a hydroxylamine derivative [VII] or its salt.

A suitable salt of a hydroxylamine derivative [VII] may be a hydrohalide (e.g. hydrochloride, etc.).

10 This reaction is usually carried out in a conventional solvent such as water, alcohol (e.g. methanol, ethanol, propanol, etc.), tetrahydrofuran, dioxane, ethyl acetate, N,N-dimethylformamide, dimethyl sulfoxide or any other organic solvent which does not adversely influence the reaction. In case the compound [VII] is liquid, it can be also used as a solvent.

The reaction temperature is not critical, and the reaction can be carried out under cooling to warming or heating.

15 The process for preparing the starting compound [II] or its salt is explained in detail in the following.

Process A

20 The compound [II'] or its salt can be prepared by halogenating a compound [VIII] or its salt.

Suitable salts of the compounds [II'] and [VIII] may be the same as those exemplified for the object compound [I].

This reaction may be carried out in substantially the same manner as Process 6, and therefore the reaction mode and reaction conditions [e.g. halogenating agent, solvent, reaction temperature, etc.] of this reaction are to be referred to those as explained in Process 6.

25 The object compounds and pharmaceutically acceptable salts thereof are novel and exhibit pharmacological activities and are useful for the treatment and prophylaxis of rheumatism (e.g. rheumatitis, etc.) nephritis, thrombocytopenia [e.g. idiopathic thrombocytopenic purpura, secondary thrombocytopenic purpura, thrombocytopenia due to a side effect of an antitumor agent (e.g. mitomycin C, etc.) etc.], tumor, side effect of an antitumor agent (e.g. decrease of body weight, etc.) and the like.

30 In order to show the utility of the object compounds [I], antirheumatic, anti-nephritic and platelet number-increasing activities and alleviating activity for side effect of antitumor agent of the object compounds [I] are explained in the following.

35

Antirheumatic activity

Test 1 Effect on collagen induced arthritis in mice

40

Method :

45 Ten male DBA/1 mice were used per group. Type II bovine collagen was dissolved in 0.1M acetic acid and emulsified in complete Freund's adjuvant (CFA). Mice were primed with 125 µg of Type II collagen in CFA intradermally at the base of the tail. Mice were challenged after 21 day with the same procedure. From the day of challenge, drug was administered orally once a day for 3 weeks and mice were inspected weekly for visual signs of arthritis. To evaluate the effect of drugs, an arthritic index was used. Arthritic index was obtained by scoring each limb 0 - 3 severity, representing joint swelling and erythema (Grade 1), visible joint disorder (Grade 2) and detectable joint ankylosis (Grade 3), and by summing up scores of four limbs.

50

55

| Results : | | |
|---------------|-----------------------|-------------------|
| Compounds | Dose level (mg/kg) | Inhibition (%) |
| a compound of | | |
| Example 18 | 100 | 50 |
| Example 23 | 100 | 51 |
| Example 27 | 100 | 46 |
| Example 29 | 100 | 57 |
| Example 30 | 100 | 35 |
| Example 33 | 100 | 44 |
| Example 37 | 100 | 50 |
| Example 38 | 100 | 31 |
| Example 40 | 100 | 44 |

20 Anti-nephritic activity

Test 2 Effect on chloric GVH disease (nephritis)

Method :

Six weeks old female (57BL/6 x DBA/2)F₁ and DBA/2 mice were used. Graft-versus-host (GVH) disease was induced in (57BL/6 x DBA/2)F₁ mice with two injections of DBA/2 spleen cells given 5 days apart. Each injection contained 5×10^7 cells. From 3 days after second cell injection, drug was administered orally once a day for 8 weeks. To assess the renal disease, at 8 weeks after last cell injection, proteinuria were measured. The concentration of serum albumin in the urine was determined by the single radial immunodiffusion method using rabbit anti-mouse serum albumin antiserum. Ten mice were used per group. Antinephritic activity of the compound was expressed the inhibition of albumin in urine.

| Results : | | |
|---------------|-----------------------|--|
| Compounds | Dose level (mg/kg) | Inhibition of albumin in urine (%) |
| a compound of | | |
| Example 7 | 100 | 52 |
| Example 18 | 100 | 98 |
| Example 23 | 100 | 96 |
| Example 29 | 100 | 90 |
| Example 30 | 100 | 98 |
| Example 33 | 100 | 70 |
| Example 35 | 100 | 74 |
| Example 37 | 100 | 100 |
| Example 38 | 100 | 78 |

Platelet number-increasing activity

Test 3 Increasing effect on the platelet number decreased by mitomycin C

Method :

5 A test compound was given orally once a day for 5 days to male ddY mice aged 6 or 7 weeks. The animals were used in groups of 10. Mitomycin C (hereinafter referred to as MMC) at a dose of 3.2 mg/kg was given intravenously to mice on day 0, 2 and 4 after the initial dosing with the test compound. The number of platelets were counted 5 days after the final dosing with the test compound, in which mice were
 10 bled from the orbital plexus and the platelets were counted with an automatic blood analyzer. The number of platelets of each group was calculated on the basis of the number of platelets (%) obtained from the non-test compound group.

15

20

25

| Results : | | |
|---------------|--------------------|-------------------------|
| Compounds | Dose level (mg/kg) | Number of platelets (%) |
| a compound of | | |
| Example 14 | 32 | 164 |
| Example 15 | 32 | 150 |
| Example 30 | 32 | 184 |
| Example 31 | 32 | 154 |
| Example 33 | 32 | 210 |
| Example 40 | 32 | 165 |
| Example 68 | 32 | 135 |

30

Alleviating activity for side effect of antitumor agent

35

Test 4 Restoring effect on the body weight decreased by MMC

Method :

40 A test compound was given orally once a day for 5 days to male ddY mice aged 6 or 7 weeks. The animals were used in groups of 10. MMC at a dose of 3.2 mg/kg was given intravenously to mice on day 0, 2 and 4 after the initial dosing with the test compound. The body weight of mice were measured on day 0 and 8.

45

The body weight of non-test compound group which was only administered MMC as above was measured on day 0 and day 8 as a control.

50

55

| Results: | | | |
|--------------------------------|-------------------------------|-----------------|-------|
| | Dose of test compound (mg/kg) | Body weight (g) | |
| | | day 0 | day 8 |
| MMC & a compound of Example 30 | 100 | 32.7 | 31.1 |
| MMC & a compound of Example 31 | 100 | 32.6 | 31.1 |
| MMC & a compound of Example 68 | 100 | 32.6 | 30.2 |
| MMC (no test compound) | - | 32.6 | 28.6 |
| (mean of 10 mice) | | | |

[I] and pharmaceutically acceptable salts thereof are used in the form of conventional pharmaceutical composition such as powders, fine granules, granules, tablets, dragee, microcapsules, capsules, suppository, solution, suspension, emulsion, syrups and the like. If desired, diluents or disintegrators (e.g. sucrose, lactose, starch, crystalline cellulose, low-substituted hydroxypropyl cellulose, synthetic aluminum silicate, etc.), binding agents (e.g. cellulose, methylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, polypropylpyrrolidone, polyvinylpyrrolidone, gelatin, gum arabic, polyethyleneglycol, etc.), coloring agents, sweetening agents, lubricant (e.g. magnesium stearate, etc.) or the like, may be dispensed with said composition.

The dosage of said composition according to this invention depends on the patient's age, body weight, condition, etc., and it is generally administered by the oral route at the daily dose level of 1 mg to 1 g as the object compound [I] or a salt thereof, preferably 10 mg to 100 mg on the same basis, at the interval of 1 to 3 times a day. Typical unit doses may be 5 mg, 10 mg, 20 mg, 50 mg, 100 mg and the like, although these are only examples and not limitative, of course.

The following Preparations and Examples are given for the purpose of illustrating the present invention in more detail.

Preparation 1

A mixture of 2-acetylamino-4-hydroxymethylthiazole (7.0 g) and N-chlorosuccinimide (6.5 g) in acetic acid (70 ml) was heated at 40 °C for 3.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was added the aqueous sodium bicarbonate.

The mixture was extracted with a mixture of ethyl acetate and tetrahydrofuran (1:1), washed with water and dried over magnesium sulfate. The solvent was concentrated under reduced pressure and the residue was triturated with isopropyl ether. The precipitates were collected by filtration, washed with isopropyl ether and dried in vacuo to give 2-acetylamino-5-chloro-4-hydroxymethylthiazole (7.3 g, yield : 78.5%).

mp : 145-146 °C

IR (Nujol) : 3150, 1690, 1550, 1285 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 2.17 (3H, s), 4.17 (2H, d, J = 5Hz), 5.17 (1H, t, J = 5Hz)

Mass : M⁺ 208, M⁺ 207, M 206, m/e 164, 147, 135

Preparation 2

A mixture of 2-amino-4-methylthiazole hydrochloride (1.5 g) and N-chlorosuccinimide (1.6 g) in acetic acid (15 ml) was heated 40 °C for 5.5 hours with stirring. The reaction mixture was poured into ice water and the solution was adjusted to pH 8.5 using sodium bicarbonate. The mixture was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give 2-amino-5-chloro-4-methylthiazole (1.4 g, yield: 94.6%, oil).

NMR (DMSO-d₆, 200MHZ, ppm) : 2.09 (3H, s), 7.00 (2H, br s)

Mass : M⁺ 150, M⁺ 149, M 148, m/e 133, 113, 99

Preparation 3

To a solution of 2-amino-4-methylthiazole hydrochloride (3.0 g) in acetic acid (20 ml) was added once N-bromosuccinimide (4.0 g) at room temperature with stirring. The mixture was stirred at room temperature for 1.5 hours and the reaction mixture was poured into isopropyl ether under ice cooling. The precipitates were collected by filtration, washed with ethyl ether and dried in vacuo to give 2-amino-5-bromo-4-methylthiazole hydrochloride (4.1 g, yield : 89.1%).

mp : 175-178° C (dec.)

IR (Nujol) : 3200, 2500-2700, 1635 cm^{-1} NMR (DMSO- d_6 , 200MHZ, ppm) : 2.14 (3H, s), 8.90 (3H, br s)

10 Mass : M^{+3} 196, M^{+2} 195, M^{+1} 194, M 193, m/e 192, 191, 149, 123, 113

Example 1

15 A solution of 1-acetyl-2-(4-nitrophenyl)ethane (9.6 g) and pyridinium bromide perbromide (18 g) in acetic acid and 35% hydrogen bromide in acetic acid (21 ml) was stirred at room temperature for 5 hours. The reaction mixture was washed with isopropyl ether and decanted. To the residue was added the thiourea (6 g), sodium acetate (8 g) and ethanol (150 ml). The mixture was heated at 50° C for 2 hours with stirring. The reaction mixture was concentrated in vacuo and to the residue was added water and then the mixture was adjusted to pH 8. The solution was extracted with ethyl acetate and washed with 10% aqueous hydrochloric acid and dried over magnesium sulfate. The solvent was concentrated in vacuo and the residue was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 300 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-4-methyl-5-(4-nitrobenzyl)thiazole (2.0 g, yield : 16.2%).

25 IR (Nujol) : 3200, 3250, 3350, 1630, 1605, 1515, 1350 cm^{-1}

NMR (DMSO- d_6 , 90MHZ, ppm) : 2.20 (3H, s), 4.10 (2H, s), 7.50 (2H, d, J=9Hz), 8.16 (2H, d, J=9Hz), 8.85 (2H, s)

Mass : M^{+1} 250, M 249, m/e 234, 204, 190, 127

30

Example 2

35 To a solution of potassium permanganate (4 g) in water (400 ml) was portionwise added the 2-amino-4-methyl-5-(4-nitrobenzyl)thiazole (2.3 g) at 80-90° C with stirring. The mixture was refluxed for 2 hours with stirring. The reaction mixture was filtered and then the filtrate was adjusted to pH 2.0 using diluted aqueous hydrochloric acid under ice cooling. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-(4-nitrobenzoyl)-4-thiazole carboxylic acid (1.50 g, yield : 75.0%).

IR (Nujol) : 3500, 2650, 2550, 1710, 1690, 1605, 1525, 1350 cm^{-1}

40 NMR (DMSO- d_6 , 60MHZ, ppm) : 7.70 (2H, s), 8.25 (2H, d, J=10Hz), 8.43 (2H, d, J=10Hz)

Mass : M 293, m/e 192, 167

Example 3

45 A mixture of 2-amino-5-(4-nitrobenzoyl)-4-thiazole carboxylic acid (2.6 g) and 10% palladium on carbon (1 g, 50% wet) in a mixture of methanol (50 ml) and tetrahydrofuran (50 ml) was hydrogenated under atmospheric pressure for 2 hours. The reaction mixture was filtered and then the filtrate was concentrated under reduced pressure. The residue was triturated with ether and the precipitates were collected by filtration, washed with ether and dried in vacuo to give 2-amino-5-(4-amino benzoyl)-4-thiazolecarboxylic acid (2.10 g, yield : 90.0%).

mp : 290-295° C (dec.)

IR (Nujol) : 3470, 3370, 2700-2500, 1690-1660, 1600 cm^{-1}

55 NMR (DMSO- d_6 , 60MHZ, ppm) : 5.73 (2H, br s), 6.80 (2H, d, J=10Hz), 6.80 (2H, br s), 7.70 (2H, d, J=10Hz)

Mass : m/e 220, 205, 151, 137, 120

Example 4

A mixture of chloromethyl-(4-nitrophenyl)diketone (2 g), thiourea (1.5 g) and sodium acetate (1.6 g) in ethanol (20 ml) was heated at 50 °C for 4 hours with stirring. The mixture was concentrated in vacuo and the residue was triturated with water. The precipitates were collected by filtration, washed with water and dried in vacuo to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 100 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-4-(4-nitrobenzoyl)thiazole (0.71 g, yield : 32.4%).

mp : 194-198 °C (dec.)

IR (Nujol) : 3300-3450, 1660, 1600, 1520, 1350 cm⁻¹

NMR (DMSO-d₆, 90MHZ, ppm) : 7.30 (2H, s), 7.60 (1H, s), 8.15 (2H, d, J=9Hz), 8.30 (2H, d, J=9Hz) Mass : M⁺ 250, M 249, m/e 219, 205, 150, 99

Example 5

A mixture of 2-amino-4-(4-nitrobenzoyl)thiazole (0.7 g) and 10% palladium on carbon (1 g, 50% wet) in a mixture of tetrahydrofuran (50 ml), methanol (50 ml) and acetic acid (5 ml) was hydrogenated under atmospheric pressure for 7 hours. The reaction mixture was filtered and then, the filtrate was concentrated under reduced pressure. The residue was dissolved in water and adjusted to pH 8.0 using aqueous sodium bicarbonate. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-4-(4-aminobenzoyl)thiazole (0.54 g, yield : 87.7%)

mp : 180-184 °C (dec.)

IR (Nujol) : 3150, 3350, 3450, 1620, 1595 cm⁻¹

NMR (DMSO-d₆, 90MHZ, ppm) : 6.00 (2H, S), 6.55 (2H, d, J=9Hz), 7.10 (2H, s), 7.25 (1H, s), 7.93 (2H, d, J=9Hz)

Mass : M⁺ 220, M 219, m/e 205, 160, 120

Example 6

A mixture of 2-amino-4-(4-aminobenzoyl)thiazole (6 g) and methoxyamine hydrochloride (13 g) in methanol (800 ml) was stirred at room temperature for 16 hours. The reaction mixture was concentrated under reduced pressure and then to the residue was added water. The solution was adjusted to pH 8.5 using 10% aqueous sodium bicarbonate under ice-cooling. The precipitates were collected by filtration, washed with water and recrystallized from ethanol to give 2-amino-4-[(4-aminophenyl)-methoxyiminomethyl]thiazole (e.9 g, yield : 72.1%)

mp : 181-183 °C

IR (Nujol) : 3350, 3100, 1605, 1510, 1380 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 3.73 (3H, s), 5.29 (2H, s), 6.46 (2H, d, J=9Hz), 6.95 (1H, s), 7.00 (2H, d, J=9Hz)

Mass : M⁺ 249, M 248, m/e 217, 203

Example 7

To a mixture of 2-amino-5-(4-nitrophenylsulfonyl)thiazole (4.0 g) and ammonium chloride in a mixture of ethanol (80 ml), tetrahydrofuran (40 ml) and water (30 ml) was portionwise added the iron powder (4 g) at 80 °C with stirring. The mixture was refluxed for 1.5 hours with stirring. The reaction mixture was filtered by suction and the filtrate was concentrated under reduced pressure. The residue was triturated with water and the precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-(4-aminophenylsulfonyl)thiazole (3.10 g, yield : 86.6%).

mp : 218-219 °C

IR (Nujol) : 3400, 3300, 1620, 1595, 1535, 1380 cm⁻¹ NMR (DMSO-d₆, 90MHZ, ppm) : 6.07 (2H, s), 6.57 (2H, d, J=9Hz), 7.43 (2H, d, J=9Hz), 7.40 (1H, s), 7.77 (2H, s)

Mass : M⁺ 256, M 255, m/e 191, 140

Example 8

A mixture of 2-acetylamino-4-chloromethylthiazole (1.9 g), 4-nitrothiophenol (1.6 g) and potassium carbonate (2.0 g) in a N,N-dimethylformamide (50 ml) was heated at 100 °C for 3 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-acetylamino-4-(4-nitrophenylthiomethyl)thiazole (2.95 g, yield : 95.5%).
 mp : 165-166 °C
 IR (Nujol) : 3150, 1655, 1595, 1545, 1500, 1335, 1290 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 2.17 (3H, s), 4.67 (2H, s), 7.15 (1H, s), 7.60 (2H, d, J=8Hz), 8.17 (2H, d, J=8Hz)
 Mass : M⁺ 310, M 309, m/e 267, 246, 155, 124, 113

Example 9

To a mixture of 2-acetylamino-4-(4-nitrophenylthiomethyl)thiazole (11 g) and ammonium chloride (2 g) in a mixture of tetrahydrofuran (200 ml), ethanol (200 ml) and water (100 ml) was added portionwise the iron powder (17 g) at 80 °C with stirring. The mixture was refluxed for 3 hours with stirring. The reaction mixture was filtered by suction and the filtrate was concentrated under reduced pressure and then the residue was triturated with water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-acetylamino-4-(4-aminophenylthiomethyl)thiazole (9.3 g, yield : 93.6%).
 IR (Nujol) : 3400, 3250, 3150, 1690, 1545, 1370, 1220 cm⁻¹
 NMR (DMSO-d₆, 90MHZ, ppm) : 2.10 (3H, s), 3.90 (2H, s), 5.20 (2H, s), 6.50 (2H, d, J=8Hz), 6.70 (1H, s), 7.05 (2H, d, J=8Hz), 12.10 (1H, s)
 Mass : M⁺ 280, M 279, m/e 236, 220, 216, 205

Example 10

To a solution of 2-acetylamino-4-(4-aminophenylthiomethyl)thiazole (9.0 g) in ethyl acetate (300 ml) was added portionwise the 3-chloroperbenzoic acid (17 g) at 5 °C with stirring. The mixture was stirred at room temperature for 16 hours. The reaction mixture was washed with aqueous sodium bicarbonate and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 300 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-acetylamino-4-(4-aminophenylsulfonylmethyl)thiazole (4.85 g, yield: 48.3%).
 mp : 135-137 °C
 IR (Nujol) : 3450, 3350, 3200, 1680, 1635, 1595, 1550, 1300 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 2.17 (3H, s), 4.50 (2H, s), 6.17 (2H, s), 6.63 (2H, d, J=8Hz), 6.90 (1H, s), 7.35 (2H, d, J=8Hz)

Example 11

A solution of 2-acetylamino-4-(4-aminophenylsulfonylmethyl)thiazole (4.8 g) in a mixture of acetic acid (35 ml) and 6N-aqueous hydrochloric acid (10 ml) was refluxed for 2.5 hours with stirring. The reaction mixture was poured into ice-water and then the solution was adjusted to pH 8.0 using 10% aqueous sodium bicarbonate with stirring. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-4-(4-aminophenylsulfonylmethyl)thiazole (2.50 g, yield : 60.2%).
 mp : 203-206 °C (dec.)
 IR (Nujol) : 3450, 3350, 1630, 1595, 1530, 1380 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 4.20 (2H, s), 6.03 (2H, s), 6.27 (1H, s), 6.57 (2H, d, J=8Hz), 6.85 (2H, s), 7.33 (2H, d, J=8Hz)
 Mass : M 269, m/e 205, 162, 140, 113

Example 12

A mixture of 4-nitrothiophenol (9.3 g), 2-amino-4-chloromethylthiazole hydrochloride (11 g) and potassium carbonate (20 g) in N,N-dimethylformamide (200 ml) was heated at 85-90 °C for 5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-4-(4-nitrophenylthiomethyl)thiazole (15.80 g, yield : 98.6%).
 IR (Nujol) : 3400, 3100, 1630, 1530, 1340 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 4.23 (2H, s), 6.60 (1H, s), 7.03 (2H, s), 7.63 (2H, d, J=9Hz), 8.20 (2H, d, J=9Hz)
 Mass : M⁺ 268, M 267, m/e 237, 177, 113

Example 13

To a mixture of 2-amino-4-(4-nitrophenylthiomethyl)thiazole (15 g) and ammonium chloride (2 g) in a mixture of tetrahydrofuran (100 ml), ethanol (150 ml) and water was added portionwise the iron powder (15 g) at 80 °C with stirring. The mixture was refluxed for 2 hours with stirring. The reaction mixture was filtered by suction and the filtrate was concentrated under reduced pressure. The residue was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with saturated aqueous sodium chloride and dried over magnesium residue was triturated with chloroform. The precipitates were collected by filtration, washed with ether and dried in vacuo to give 2-amino-4-(4-aminophenylthiomethyl)thiazole (10.50 g, yield : 73.0%)
 mp : 130-132 °C
 IR (Nujol) : 3425, 3350, 1630, 1605, 1595, 1535, 1495, 1440, 1380, 1340, 1280 cm⁻¹
 NMR (DMSO-d₆, 90MHZ, ppm) : 3.70 (2H, s), 5.15 (2H, s), 6.10 (1H, s), 6.45 (2H, d, J=9Hz), 6.83 (2H, s), 7.00 (2H, d, J=9Hz)
 Mass : M⁺ 238, M 237, m/e 204, 124, 113

Example 14

A solution of 3-chloroperbenzoic acid (4.9 g) in dichloromethane (100 ml) was dropwise added to a solution of 2-amino-4-(4-aminophenylthiomethyl)thiazole (5.1 g) in a mixture of dichloromethane (200 ml) and N,N-dimethylformamide (10 ml) at 5 °C with stirring. The mixture was stirred at 5 °C for 1.5 hours with stirring. The precipitates were collected by filtration, washed with ethyl acetate and dried in vacuo to give solid. The solid was recrystallized from ethanol to give 2-amino-4-(4-aminophenylsulfinylmethyl)thiazole (4.70 g, yield : 86.3%).
 IR (Nujol) : 3350-3100, 1620, 1600, 1500, 1380, 1300 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 3.87 (2H, s), 6.27 (2H, s), 6.67 (2H, d, J=9Hz), 7.00 (2H, s), 7.30 (2H, d, J=9Hz), 7.67 (1H, s)
 Mass : M⁺ 254, M 253, m/e 237, 205, 156, 139

Example 15

To a solution of 2-amino-4-(4-aminophenylsulfinylmethyl)thiazole (2.8 g) in N,N-dimethylformamide (30 ml) was added portionwise the 3-chloroperbenzoic acid (2.6 g) at 5 °C with stirring. The mixture was stirred at room temperature for 2 hours and then the solution was poured into ice-water. The precipitates were collected by filtration, washed with aqueous sodium bicarbonate, washed with water and dried in vacuo to give 2-amino-4-(4-aminophenylsulfonylmethyl)thiazole (2.85 g, yield 95.6%).
 mp : 204-208 °C (dec.)
 IR (Nujol) : 3375, 3275, 3150, 1615, 1595, 1295, 1140 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 4.30 (2H, s), 6.10 (2H, s), 6.30 (1H, s), 6.67 (2H, d, J=8Hz), 6.95 (2H, s), 7.43 (2H, d, J=8Hz)
 Mass : M 269, m/e 220, 205

Example 16

A mixture of 2-acetylamino-5-chlorothiazole (5 g), 4-nitrothiophenol (4.83 g) and potassium carbonate (7.8 g) in N,N-dimethylformamide (100 ml) was heated at 120 °C for 3 hours with stirring. The reaction mixture was poured into ice-water. The precipitates were collected by filtration, washed with water and dried in vacuo to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 200 g) and eluted with a mixture of n-hexane and ethyl acetate (3:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-acetylamino-5-(4-nitrophenylthio)thiazole (3.74 g, yield : 50.2%).

mp : 250-255 °C (dec.)
 IR (Nujol) : 3150, 1695, 1595, 1550, 1505, 1340, 1300, 1230 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 2.23 (3H, s), 7.40 (2H, d, J=8Hz), 7.90 (1H, s), 8.23 (2H, d, J=8Hz), 12.43 (1H, br s)
 Mass : M⁺ 296, M 295, m/e 265, 253, 223, 181, 166

Example 17

To a mixture of 2-acetylamino-5-(4-nitrophenylthio)thiazole (2.8 g) and ammonium chloride (0.3 g) in a mixture of ethanol (60 ml), water (30 ml) and tetrahydrofuran (20 ml) was portionwise added the iron powder (3 g) at 80 °C with stirring. The mixture was refluxed for 2.5 hours with stirring. The reaction mixture was filtered by suction and the filtrate was concentrated under reduced pressure. The residue was triturated with water, the precipitates were collected by filtration, washed with water and dried in vacuo to give 2-acetylamino-5-(4-aminophenylthio)thiazole (2.0 g, yield : 79.5%).

mp : 255-257 °C (dec.)
 IR (Nujol) : 3450, 3375, 1680, 1620, 1595, 1380, 1300 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 2.17 (3H, s), 5.40 (2H, s), 6.60 (2H, d, J=9Hz), 7.20 (2H, d, J=9Hz), 7.58 (1H, s), 12.15 (1H, s)
 Mass : M⁺ 266, M 265, m/e 223, 191, 181

Example 18

A solution of 2-acetylamino-5-(4-aminophenylthio)thiazole (2.5 g) in a mixture of acetic acid (20 ml) and 6N-aqueous hydrochloric acid (5 ml) was refluxed for 4 hours with stirring. The reaction mixture was poured into ice-water and the solution was adjusted to pH 10 using 1N-aqueous sodium hydroxide with stirring under ice cooling. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-(4-aminophenylthio)thiazole (1.70 g, yield : 81.5%).

IR (Nujol) : 3400, 3300, 3150, 1630, 1600, 1515, 1380 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 5.27 (2H, s), 6.60 (2H, d, J=9Hz), 7.10 (2H, d, J=9Hz), 7.17 (1H, s), 7.27 (2H, s)
 Mass : M⁺ 224, M 223, m/e 191, 164, 136, 125

Example 19

To a solution of 2-amino-5-(4-aminophenylthio)thiazole (4.0 g) in a mixture of chloroform (140 ml) and N,N-dimethylformamide (20 ml) was dropwise added the solution of 3-chloroperbenzoic acid (4.65 g) in chloroform (50 ml) at 5 °C with stirring. The mixture was stirred at 5 °C for 4 hours. The precipitates were collected by filtration, washed with 10% aqueous sodium bicarbonate and water. The solid was dried in vacuo to give 2-amino-5-(4-aminophenylsulfinyl)thiazole (3.75 g, yield : 87.5%).

mp : 173-175 °C
 IR (Nujol) : 3350, 3350, 3225, 1640, 1595, 1525, 1380 1320, 1225 cm⁻¹
 NMR (DMSO-d₆, 90MHZ, ppm) : 5.67 (2H, s), 6.97 (2H, d, J=9Hz), 7.23 (2H, d, J=9Hz), 7.50 (1H, s), 7.62 (2H, s)
 Mass : M⁺ 240, M 239, m/e 223, 191, 147, 140

Example 20

To a solution of 2-acetylamino-5-(4-aminophenylthio)thiazole (6.6 g) in a mixture of dichloromethane (300 ml) and N,N-dimethylformamide (50 ml) was dropwise added the solution of 3-chloroperbenzoic acid (5.9 g) in dichloromethane (100 ml) at 5 °C with stirring. The mixture was stirred at 5 °C for 2 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was poured into 10% aqueous sodium bicarbonate. The solution was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with saturated aqueous sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give 2-acetylamino-5-(4-aminophenylsulfinyl)thiazole (5.0 g, yield : 71.1%).
 mp : 194-196 °C (dec.)
 IR (Nujol) : 3350, 3175, 1710, 1630, 1695, 1550, 1380, 1300, 1230 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 2.17 (3H, s), 5.80 (2H, br s), 6.70 (2H, d, J=9Hz), 7.37 (2H, d, J=9Hz), 7.9 (1H, s)
 Mass : M⁺ 282, M 281, m/e 265, 234, 222, 191

Example 21

A mixture of 2-acetylamino-5-(4-aminophenylsulfinyl)thiazole (5.0 g) in a mixture of aqueous 6N-hydrochloric acid (10 ml) and acetic acid (35 ml) was refluxed for 3.5 hours with stirring. The reaction mixture was diluted with water and adjusted to pH 8 using aqueous sodium carbonate under ice cooling. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-(4-aminophenylsulfinyl)thiazole (4.5 g, yield : 100%).
 mp : 205-208 °C (dec.)
 NMR (DMSO-d₆, 60MHZ, ppm) : 5.27 (2H, s), 6.55 (2H, d, J=8Hz), 6.67 (2H, d, J=8Hz), 7.17 (1H, s), 7.42 (2H, s)
 Mass : m/e 223, 191, 124, 99

Example 22

A mixture of 2-acetylamino-5-chlorothiazole (5.3 g), 4-mercaptopyridine (3.4 g) and potassium carbonate in N,N-dimethylformamide (50 ml) was heated at 120 °C for 2.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-acetylamino-5-(4-pyridylthio)thiazole (6.3 g, yield : 83.7%).
 IR (Nujol) : 3150, 1680, 1580, 1300 cm⁻¹
 NMR (DMSO-d₆, 90MHZ, ppm) : 2.23 (3H, s), 7.10 (2H, d, J=6Hz), 7.80 (1H, s), 8.40 (2H, d, J=6Hz), 11.90 (1H, br s)

Example 23

A mixture of 2-acetylamino-5-(4-pyridylthio)thiazole (4.7 g), acetic acid (35 ml) and 6N-aqueous hydrochloric acid (10 ml) was refluxed for 2 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted pH 8.5 using aqueous sodium bicarbonate under ice cooling. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-(4-pyridylthio)thiazole (2.7 g, yield : 69.5%).
 mp : 180-185 °C (dec.)
 IR (Nujol) : 3270, 3150, 1630, 1580, 1380 cm⁻¹
 NMR (DMSO-d₆, 90MHZ, ppm) : 7.13 (2H, d, J=6Hz), 7.30 (1H, s), 7.60 (2H, s), 8.40 (2H, d, J=6Hz)
 Mass : M⁺ 210, M 209, m/e 188, 150, 131, 99

Example 24

To a mixture of 2-amino-5-(4-pyridylthio)thiazole (4.0 g) in a mixture of chloroform (300 ml) and N,N-

dimethylformamide (10 ml) was added dropwise the solution of 3-chloroperbenzoic acid (4.5 g) in chloroform (100 ml) at 5 °C with stirring. The mixture was stirred at 5 °C for 26 hours under ice cooling. The reaction mixture was washed with aqueous sodium bicarbonate and water and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subject to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 250 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-5-(4-pyridylsulfinyl)thiazole (2.5 g, yield: 58.1%)
 mp : 193-195 °C
 IR (Nujol) : 3350, 3250, 1610, 1575, 1525, 1280, 1220 cm⁻¹
 10 NMR (DMSO-d₆, 60MHZ, ppm) : 7.62 (2H, d, J=6Hz), 7.87 (1H, s), 7.97 (2H, s), 8.80 (2H, d, J=6Hz)
 Mass : M 255, m/e 209, 177, 147, 131

Example 25

15 A mixture of 2-acetylamino-5-(4-nitrophenylthio)thiazole (4.0 g) in a mixture of acetic acid (30 ml) and aqueous 6N hydrochloric acid (9 ml) was refluxed for 3 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 8.5 using aqueous sodium bicarbonate. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-(4-nitrophenylthio)thiazole (2.6 g, yield : 76.5%).
 mp : 162-164 °C
 IR (Nujol) : 3420, 3270, 1680, 1595, 1580, 1530, 1335, 1215 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 7.30 (2H, d, J=8Hz), 7.40 (1H, s), 7.50 (2H, s), 8.20 (2H, d, J=8Hz) Mass: M⁺ 254, M 253, m/e 223, 191, 164, 149, 121, 99

Example 26

25 To a mixture of 2-amino-5-(4-nitrophenylthio)thiazole (2.6 g) and pyridine (1 g) in N,N-dimethylformamide (30 ml) was added dropwise the propionylchloride (1.1 g) at 5 °C under ice cooling with stirring. The mixture was stirred at 5 °C for 3.5 hours. The reaction mixture was poured into ice water and the mixture was adjusted to pH 8 using aqueous sodium bicarbonate. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-propionylamino-5-(4-nitrophenylthio)thiazole (2.5 g, yield : 78.7%).
 35 mp : 227-229 °C (dec.)
 IR (Nujol) : 3150, 1710, 1595, 1580, 1555, 1505, 1340, 1180 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 1.20 (3H, t, J=8Hz), 2.6 (2H, g, J=8Hz), 7.45 (2H, d, J=8Hz), 7.92 (1H, s), 8.23 (2H, d, J=8Hz)
 Mass : M⁺ 310, M 309, m/e 280, 252, 222

Example 27

45 To a mixture of 2-propionylamino-5-(4-nitrophenylthio)thiazole (3 g), and ammonium chloride (1 g) in a mixture of ethanol (100 ml), water (30 ml) and tetrahydrofuran (70 ml) was portionwise added the iron powder at 80 °C with stirring. The mixture was refluxed for 2 hours with stirring. The reaction mixture was filtered by suction and the filtrate was concentrated under reduced pressure. The residue was triturated with water, the precipitates were collected by filtration and washed with water to give solid. The solid was recrystallized from ethanol to give 2-propionylamino-5-(4-aminophenylthio)thiazole (2.6 g, yield : 96.3%).
 50 mp : 185-187 °C
 NMR (DMSO-d₆, 60MHZ, ppm) : 1.30 (3H, t, J=7Hz), 2.50 (2H, q, J=7Hz), 5.50 (2H, br s), 6.60 (2H, d, J=8Hz), 7.23 (2H, d, J=8Hz), 7.60 (1H, s)
 Mass : M⁺ 280, M 279, m/e 264, 250, 222, 205

Example 28

A mixture of 2-acetylamino-5-chlorothiazole (5.3 g), 2-mercaptopyridine (3.5 g) and potassium carbonate

(6.2 g) in N,N-dimethylformamide (50 ml) was heated at 130 °C for 3.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-acetylthio-5-(2-pyridylthio)thiazole (5.70 g, yield : 76.0%).

5 mp : 185-188 °C (dec.)

IR (Nujol) : 3150, 1695, 1575, 1300, 1280, 1230 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 2.20 (3H, s), 7.00-7.40 (2H, m), 7.70-7.90 (2H, m), 8.50 (1H, m), 12.40 (1H, s)

Mass : M⁺ 252, M 251, m/e 209, 176, 167

10

Example 29

A mixture of 2-acetylthio-5-(2-pyridylthio)thiazole (5.0 g) in a mixture of acetic acid (50 ml) and aqueous 6N-hydrochloric acid (10 ml) was refluxed for 2 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 8.5 using aqueous sodium bicarbonate, and then the mixture was extracted with ethyl acetate. The organic layer was washed with water and dried over magnesium sulfate. The solvent was concentrated under reduced pressure and the residue was triturated with a solution of hydrochloric acid in ethanol. The precipitates were collected by filtration, washed with isopropyl ether and dried in vacuo to give 2-amino-5-(2-pyridylthio)thiazole dihydrochloride (4.60 g, yield : 85.8%).

mp : 220-225 °C (dec.)

IR (Nujol) : 2550-2300, 1620, 1595 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 7.20-7.60 (2H, m), 7.70-8.00 (4H, m), 8.55 (1H, m), 10.50 (3H, br s)

25 Mass : m/e 209 (free), 187, 167, 123

Example 30

30 To a solution of 2-amino-5-(2-pyridylthio)thiazole dihydrochloride (4.0 g) in chloroform (100 ml) was dropwise added the solution of 3-chloroperbenzoic acid (5.0 g) in chloroform (100 ml) at 5 °C with stirring. The mixture was stirred at 5 °C for 1.5 hours. The reaction mixture was washed with aqueous sodium bicarbonate and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 100 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-5-(2-pyridylsulfonyl)thiazole (3.4 g, yield : 78.9%).

mp : 200-202 °C (dec.)

IR (Nujol) : 3300, 3150, 1630, 1575, 1270, 1225 cm⁻¹

40 NMR (DMSO-d₆, 90MHZ, ppm) : 7.40-7.60 (1H, m), 7.70 (1H, s), 7.73 (2H, s), 7.90-8.20 (2H, m), 8.60 (1H, m)

Mass : M 225, m/e 209, 147, 115

45 Example 31

To a solution of 2-amino-5-(2-pyridylthio)thiazole (5.7 g) in chloroform (450 ml) was dropwise added the solution of 3-chloroperbenzoic acid (15 g) in chloroform (100 ml) at 5 °C with stirring. The mixture was stirred at room temperature for 16 hours. The reaction mixture was washed with aqueous sodium bicarbonate and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give 2-amino-5-(2-pyridylsulfonyl)thiazole (2.2 g, yield : 33%).

mp : 178-182 °C (dec.)

IR (Nujol) : 3375, 3300, 3150, 1645, 1610, 1525, 1320, 1220 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 7.55-7.80 (2H, m), 8.00-8.30 (4H, m), 8.77 (1H, m)

55 Mass : M⁺ 242, M 241, m/e 177, 156, 135

Example 32

- A mixture of 2-acetamido-5-chlorothiazole (14.3 g), 2-mercaptopyrimidine (10 g) and potassium carbonate anhydrous (22.4 g) in N,N-dimethylformamide (280 ml) was stirred at 150 °C an hour. The reaction mixture was poured into water with stirring under ice cooling. The mixture was extracted with ethyl acetate, washed with water and dried over magnesium sulfate. The organic layer was concentrated under reduced pressure to give solid. The solid was triturated with water, and the precipitates were collected by filtration, washed with water and dried in vacuo to give 2-acetamido-5-(2-pyrimidinylthio)thiazole (12.30 g, yield : 60.2%).
- mp : 225 °C (dec.)
 IR (Nujol) : 3170, 1645, 1555, 1310 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 2.20 (3H, s), 7.33 (1H, t, J=4Hz), 7.70 (1H, s), 8.68 (2H, d, J=4Hz), 12.33 (1H, br s)
 Mass : M⁺ 254, M⁺ 253, M 252, m/e 210, 168

Example 33

- Starting from 2-acetamido-5-(2-pyrimidinylthio)thiazole, 2-amino-5-(2-pyrimidinylthio)thiazole (2.02 g, yield : 22.0%) was obtained according to a similar manner to that of Example 40.
- mp : 175-177 °C
 IR (Nujol) : 3270, 3100, 1655, 1565, 1555, 1540 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 7.13-7.57 (4H, m), 8.40-8.77 (2H, m)
 Mass : M⁺ 212, M⁺ 211, M 210, m/e 168, 124

Example 34

- A mixture of 2-acetyl-amino-5-chlorothiazole (5.3 g), 1-methyl-2-mercaptoimidazole (3.6 g) and potassium carbonate (6.2 g) in N,N-dimethylformamide (50 ml) was heated at 130 °C for 5.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The precipitates were collected by filtration washed with water and dried in vacuo to give 2-acetyl-amino-5-(1-methylimidazol-2-ylthio)thiazole (6.95 g, yield : 91.2%).
- mp : 155-160 °C (dec.)
 IR (Nujol) : 3400, 1690, 1565, 1300 cm⁻¹ NMR (DMSO-d₆, 90MHZ, ppm) : 2.10 (3H, s), 3.70 (3H, s), 6.90 (1H, s), 7.26 (1H, s), 7.60 (1H, s)
 Mass : M⁺ 255, M 254, m/e 212, 179, 170, 114

Example 35

- A solution of 2-acetyl-amino-5-(1-methylimidazol-2-ylthio)thiazole (7.0 g) in a mixture of acetic acid (100 ml) and aqueous 6N-hydrochloric acid (20 ml) was refluxed for 3.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was adjusted to pH 8 using aqueous sodium bicarbonate under ice cooling. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-(1-methylimidazol-2-ylthio)thiazole (4.9 g, yield : 83.9%).
- mp : 180-190 °C (dec.)
 IR (Nujol) : 3300, 3150, 1620, 1530, 1280, 1220 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 3.77 (3H, s), 7.03 (1H, s), 7.25 (1H, s), 7.37 (1H, s)
 Mass : M⁺ 213, M 212, m/e 179, 170, 126, 114

Example 36

- A solution of 2-acetamido-5-(4-aminophenylthio)thiazole (3.2 g) in pyridine (64 ml) was added methanesulfonyl chloride (1.52 g) at 5 °C with stirring. The reaction mixture was stirred for 3 hours at 5 °C and concentrated in vacuo to give solid. The solid was subject to column chromatography on silica gel (silica gel, 70-230 mesh; Merck : 200 g) and eluted with a mixture of chloroform and methanol (50:1) and eluted with a mixture of chloroform and methanol (20:1). The fractions containing the objective compound were combined and evaporated to dryness in vacuo to give 2-acetamido-5-(4-methanesulfonylamino)phenyl-

thio)thiazole (4.0 g, yield : 96.6%).

mp : 236-239 °C

IR (Nujol) : 3250, 3150, 1695, 1565, 1495, 1330 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 2.16 (3H, s), 3.30 (3H, s), 7.20-7.30 (5H, m), 7.73 (1H, s), 8.10 (1H, s)

5 Mass : M⁺ 343, m/e 342, 301, 264, 222

Example 37

10 Starting from 2-acetamido-5-(4-methanesulfonylaminothiophenylthio)thiazole, 2-amino-5-(4-methanesulfonylaminothiophenylthio)thiazole (2.28 g, yield : 64.9%) was obtained according to a similar manner to that of Example 40.

mp : 185-187 °C

IR (Nujol) : 3430, 3260, 1610, 1510, 1320 cm⁻¹

15 NMR (DMSO-d₆, 60MHZ, ppm) : 3.00 (3H, s), 7.20-7.37 (5H, m), 7.47 (2H, br s), 9.76 (1H, br s)

Mass : M⁺ 302, M⁺ 301, m/e 222, 190

Example 38

20 To a mixture of 2-amino-5-(4-pyridylthio)thiazole (2.5 g) and pyridine (3 g) in N,N-dimethylformamide (25 ml) was added dropwise the 4-fluorobenzoyl chloride (2.7 g) at 5 °C under ice cooling with stirring. The mixture was stirred at 5 °C for 4 hours under ice cooling. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The precipitates were collected by filtration,
25 washed with water and dried in vacuo to give 2-(4-fluorobenzoylamino)-5-(4-pyridylthio)thiazole (2.5 g, yield : 63.1%)

mp : 220-225 °C (dec.)

IR (Nujol) : 3150, 1670, 1605, 1587, 1550, 1295, 1230 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 7.10-7.67 (4H, m), 7.95 (1H, s), 8.10-8.60 (4H, m), 12.85 (1H, s)

30 Mass : M⁺ 332, M⁺ 331, m/e 209, 123, 95,

Example 39

35 A mixture of salt of potassium 2,4-difluorothiophenol (20 g), 2-acetamido-5-chlorothiazole (21 g) and potassium carbonate anhydrous (29.8 g) in N,N-dimethylformamide (400. ml) was stirred at 130 °C for 7 hours. The reaction mixture was concentrated under reduced pressure. The residue was triturated with water and the precipitates were collected by filtration, washed with water and dried in vacuo to give solid. The solid was subjected to column chromatography on silica gel (silica gel, 70-230 mesh; Merck : 750 g)
40 and eluted with a mixture of chloroform and methanol (50:1). The fractions containing the objective compound were combined and evaporated to dryness in vacuo to give 2-acetamido-5-(2,4-difluorophenylthio)thiazole (11.91 g, yield : 38.5%). mp : 156-170 °C (dec.)

IR (Nujol) : 3160, 3060, 1695, 1585, 1560, 1295 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 2.20 (3H, s), 6.90-7.90 (4H, m), 12.27 (1H, br s)

45 Mass : M⁺ 286, m/e 270, 243

Example 40

50 A mixture of 2-acetamido-5-(2,4-difluorophenylthio)thiazole (14.8 g) in a mixture of ethanol (150 ml) and concentrated hydrochloric acid (15 ml) was refluxed for 1.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 12 using aqueous sodium hydroxide with stirring under ice-cooling. The mixture was extracted with ethyl acetate, washed with water and dried over magnesium sulfate. The organic layer was concentrated under
55 reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel, 230-400 mesh, Nakarai) (300 g) and eluted with a mixture of chloroform and methanol (100:1). The fractions containing the objective compound were combined and evaporated to dryness in vacuo to give 2-amino-5-(2,4-difluorophenylthio)thiazole (6.26 g, yield : 49.6%).

mp : 116-117° C

IR (Nujol) : 3410, 3090, 1625, 1600, 1515 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 7.07-7.76 (6H, m) MASS : M⁺ 246, M 244, m/e 157

5

Example 41

Starting from 2-amino-5-(2,4-difluorophenylthio)thiazole, 2-amino-5-(2,4-difluorophenylsulfinyl)thiazole (2.37 g, yield : 65.4%) was obtained according to a similar manner to that of Example 30.

10 mp : 171-172° C

IR (Nujol) : 3300, 3100, 1635, 1605, 1600 cm⁻¹

NMR (DMSO-d₆, 90MHZ, ppm) : 7.23-8.00 (6H, m) Mass : M⁺ 260, m/e 244, 212

15

Example 42

A mixture of 4-chloromethyl-2-formylaminothiazole (1.86 g), 4-mercaptopyridine (1.23 g) and potassium carbonate (1.8 g) in N,N-dimethylformamide (20 ml) was heated at 100° C for 2 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-formylamino-4-(4-pyridylthiomethyl)thiazole (1.7 g, yield : 68.0%).

20 mp : 182-184° C

IR (Nujol) : 1675, 1650, 1585, 1560, 1270 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 4.30 (2H, s), 7.15 (1H, s), 7.33 (2H, d, J=6Hz), 8.33 (2H, d, J=6Hz), 8.45 (1H, s)

25

Mass : M⁺ 252, M 251, m/e 223, 155, 141, 113

Example 43

30

A mixture of 2-formylamino-4-(4-pyridylthiomethyl)thiazole (1.6 g) and N-chlorosuccinimide (1.5 g) in acetic acid (25 ml) was heated at 40-50° C for 5 hours with stirring and then the mixture was allowed to stand at room temperature for 16 hours. The reaction mixture was concentrated under reduced pressure and the residue was triturated with aqueous sodium bicarbonate. The precipitates were collected by filtration washed with water and dried in vacuo to give 5-chloro-2-formylamino-4-(4-pyridylthiomethyl)thiazole (0.85 g, yield : 46.8%).

35

mp : 200-203° C (dec.)

IR (Nujol) : 1680, 1665, 1587, 1300 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 4.37 (2H, s), 7.40 (2H, d, J=6Hz), 7.47 (2H, d, J=6Hz), 8.53 (1H, s), 12.50 (1H, s)

40

Mass : M⁺ 288, M⁺ 287, M⁺ 286, M 285, m/e 256, 250, 175, 147

Example 44

45

A solution of 5-chloro-2-formylamino-4-(4-pyridylthiomethyl)thiazole (4.9 g) in a mixture of ethanol (25 ml), tetrahydrofuran (20 ml) and concentrated hydrochloric acid (7 ml) was stirred at room temperature for 4 hours. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 8 using aqueous sodium bicarbonate under ice-cooling. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-chloro-4-(4-pyridylthiomethyl)thiazole (0.26 g, yield : 56.6%).

50

IR (Nujol) : 3350, 3250, 1685, 1530 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 4.13 (2H, s), 7.33 (2H, d, J=6Hz), 7.43 (2H, s), 8.50 (2H, d, J=6Hz)

Mass : M⁺ 259, M⁺ 258, M 257, m/e 220, 147, 111

55

Example 45

A mixture of 4-chloromethyl-2-formylaminothiazole (1.76 g), 4-nitrothiophenol (1.7 g) and potassium carbonate (1.8 g) in N,N-dimethylformamide (20 ml) was heated at 100 °C with stirring. The reaction mixture was poured into ice-water and stirred at 5 °C for an hour. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-formylamino-4-(4-nitrophenylthiomethyl)thiazole (2.3 g, yield : 78%).

mp : 158-160 °C

IR (Nujol) : 3500, 1680, 1595, 1550, 1330 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 4.40 (2H, s), 7.13 (1H, s), 7.56 (2H, d, J=8Hz), 8.10 (2H, d, J=8Hz), 8.50 (1H, s)

Mass : M 295, m/e 265, 141, 113

Example 46

To a mixture of 2-formylamino-4-(4-nitrophenylthiomethyl)thiazole (2.2 g) and ammonium chloride (0.5 g) in a mixture of tetrahydrofuran (30 ml), ethanol (50 ml) and water (10 ml) was added portionwise the iron powder at 80 °C with stirring. The mixture was refluxed for 2 hours with stirring. The reaction mixture was filtered by suction and the residue was triturated with water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-formylamino-4-(4-aminophenylthiomethyl)thiazole (1.6 g, yield : 81%).

mp : 180-182 °C

IR (Nujol) : 3350, 3300, 1680, 1625, 1600, 1325, 1290 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 4.00 (2H, s), 5.23 (2H, s), 6.57 (2H, d, J=8Hz), 6.83 (1H, s), 7.10 (2H, d, J=8Hz), 8.50 (1H, s)

Mass : M⁺ 266, M 265, m/e 237, 205, 141, 124

Example 47

A mixture of acetic anhydride (1.84 g) and formic acid (0.9 g) was heated at 50 °C for 0.5 hours with stirring. The solution was cooled at room temperature and to the solution was added the 2-formylamino-4-(4-aminophenylthiomethyl)thiazole (1.6 g). The mixture was stirred at room temperature for 6.5 hours and then the mixture was poured into ice-water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-formylamino-4-(4-formylaminophenylthiomethyl)thiazole (1.7 g, yield : 96.7%).

mp : 195-197 °C (dec.)

IR (Nujol) : 3150, 1680, 1660, 1595, 1525, 1310, 1290 cm⁻¹

NMR (DMSO-d₆, 60MHZ, ppm) : 4.30 (2H, s), 7.10 (1H, s), 7.47 (2H, d, J=8Hz), 7.73 (2H, d, J=8Hz), 8.40 (1H, s), 8.60 (1H, s), 10.33 (1H, s), 12.20 (1H, s)

Mass : M⁺ 294, M 293, m/e 265, 153, 141, 113

Example 48

To a solution of 2-formylamino-4-(4-formylaminophenylthiomethyl)thiazole (2.9 g) in acetic acid (30 ml) was portionwise added N-chlorosuccinimide at 50 °C with stirring. The mixture was heated at 50 °C with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The mixture was extracted with a mixture of ethyl acetate and tetrahydrofuran (1:1), washed with water and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 150 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 5-chloro-2-formylamino-4-(4-formylaminophenylthiomethyl)thiazole (2.0 g, yield : 61.2%).

mp : 130-150 °C (dec.)

IR (Nujol) : 3350, 3200, 1710, 1690-1640, 1590 cm⁻¹

NMR (DMSO-d₆, 90MHZ, ppm) : 4.80 (2H, s), 7.26 (2H, d, J=8Hz), 7.56 (2H, d, J=8Hz), 8.25 (1H, s), 8.50 (1H, s), 10.23 (1H, s), 12.57 (1H, s)

Mass : M 327, m/e 298, 292, 263, 234, 204

Example 49

A solution of 5-chloro-2-formylamino-4-(4-formylaminophenylthiomethyl)thiazole (3.5 g) in a mixture of concentrated hydrochloric acid (9 ml), methanol (30 ml) and tetrahydrofuran (30 ml) was stirred at room temperature for 4 hours. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 8 using aqueous sodium bicarbonate with stirring under ice cooling. The precipitates were collected by filtration, washed with water and dried in vacuo to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 150 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-4-(4-aminophenylthiomethyl)-5-chlorothiazole (2.3 g, yield : 79.3%).

mp : 158-163 °C (dec.)
 IR (Nujol) : 3325, 3200, 3150, 1620, 1595, 1495, 1325, 1290 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 4.33 (2H, s), 5.50 (2H, br s), 6.60 (2H, d, J=8Hz), 7.10 (2H, d, J=8Hz), 7.25 (2H, br s)
 Mass : M 271, m/e 267, 236, 221, 204, 124

Example 50

A mixture of 2-acetylamino-5-chloro-4-hydroxymethylthiazole (1 g), 4-mercaptopyridine (0.6 g) and potassium carbonate (1 g) in N,N-dimethylformamide (20 ml) was heated at 110 °C for 8 hours with stirring. The reaction mixture was poured into ice water and filtered by suction. The filtrate was extracted with a mixture of ethyl acetate and tetrahydrofuran (1:1) and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 75 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-acetylamino-4-hydroxymethyl-5-(4-pyridylthio)thiazole (0.90 g, yield : 64.3%).

mp : 220-222 °C (dec.)
 NMR (DMSO-d₆, 90MHZ, ppm) : 2.16 (3H, s), 4.40 (2H, d, J=6Hz), 5.13 (1H, t, J=6Hz), 7.05 (2H, d, J=6Hz), 8.30 (2H, d, J=6Hz), 12.43 (1H, s)
 Mass : M⁺ 282, M 281, m/e 239, 220, 205, 188

Example 51

A mixture of 2-acetylamino-4-hydroxymethyl-5-(4-pyridylthio)thiazole (3.0 g) in a mixture of concentrated hydrochloric acid (8 ml) and ethanol (100 ml) was refluxed for 2.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with acetone. The precipitates were collected by filtration, washed with isopropyl ether and were recrystallized from a mixture of ethanol and isopropyl ether to give 2-amino-4-hydroxymethyl-5-(4-pyridylthio)thiazole dihydrochloride (2.5 g, yield : 75.3%).

mp : 231-237 °C (dec.)
 IR (Nujol) : 3350, 2300, 1610, 1560 cm⁻¹
 NMR (DMSO-d₆, 90MHZ, ppm) : 4.30-4.55 (3H, m), 7.80 (2H, d, J=6Hz), 8.65 (2H, d, J=6Hz), 8.83 (4H, br s)
 Mass : M 239, m/e 222, 210, 188

Example 52

Starting from 2-amino-5-(4-pyridylthio)thiazole 2-amino-5-(4-pyridylsulfonyl)thiazole (0.73 g, yield : 17.1%) was obtained according to a similar manner to that of Example 31.

mp : 217 °C (dec.)
 IR (Nujol) : 3260, 3100, 1620, 1580, 1525 cm⁻¹
 NMR (DMSO-d₆, 90MHZ, ppm) : 7.73-7.86 (3H, m), 8.20 (2H, br s), 8.55 (2H, d, J=6Hz)
 Mass : M⁺ 243, M⁺ 242, M+ 241, m/e 209, 195

Example 53

- A mixture of 2-amino-5-(4-aminophenylthio)thiazole, (4.0 g) in a mixture of acetic acid (40 ml) and acetic anhydride (2.2 g) was stirred at room temperature for 4 hours. The reaction mixture was concentrated under reduced pressure and the residue was triturated with aqueous sodium bicarbonate. The precipitates were collected by filtration, washed with water and dried in vacuo to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 150 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-5-(4-acetylaminophenylthio)thiazole (2.1 g, yield : 44.2%).
- mp : 240-245 °C (dec.)
 IR (Nujol) : 3400, 3325, 3200, 1660, 1605, 1595, 1320 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 2.10 (3H, s), 7.26 (2H, d, J=8Hz), 7.30 (1H, s), 7.50 (2H, s), 7.67 (2H, d, J=8Hz), 10.03 (1H, s)
- Mass : M⁺ 266, M 265, m/e 223, 207, 190

Example 54

- To a mixture of 2-amino-5-(4-methanesulfonylaminophenylthio)thiazole (2.0 g) in chloroform (100 ml) was added dropwise the solution of 3-chloroperbenzoic acid (1.6 g) in chloroform (50 ml) at 5 °C under ice cooling with stirring. The mixture was stirred at 5 °C for 2.5 hours. The reaction mixture was washed with aqueous sodium bicarbonate and the precipitates were collected by filtration. The solid was washed with aqueous sodium bicarbonate and water, dried in vacuo to give 2-amino-5-(4-methanesulfonylaminophenylsulfinyl)thiazole (1.95 g, yield : 92.6%).
- mp : 201-203 °C (dec.)
 IR (Nujol) : 3320, 3250, 3100, 1615, 1515, 1325, 1220, 1150 cm⁻¹
 NMR (DMSO-d₆, 60MHZ, ppm) : 3.17 (3H, s), 7.43 (2H, d, J=8Hz), 7.67 (2H, d, J=8Hz), 7.73 (1H, s), 7.83 (2H, s), 10.23 (1H, s)
- Mass : m/e 301, 222, 190, 146, 124, 100

Example 55

- A mixture of 2-acetylamino-5-chlorothiazole (1.76 g), 4-hydroxythiophenol (1.3 g) and potassium carbonate (2 g) in N,N-dimethylformamide (30 ml) was heated at 120 °C for 2.5 hours with stirring. The reaction mixture was poured into ice water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-acetylamino-5-(4-hydroxyphenylthio)thiazole (1.5 g, yield : 56.6%).
- mp : 265-267 °C
 IR (Nujol) : 3300, 3200, 1675, 1600, 1570, 1305, 1260 cm⁻¹
 NMR (DMSO-d₆, 200MHZ, ppm) : 2.14 (3H, s), 6.75 (2H, d, J=9Hz), 7.20 (2H, d, J=9Hz), 7.63 (1H, s), 9.69 (1H, s), 12.28 (1H, s)
- Mass : M⁺ 267, M 266, m/e 224, 191, 182, 165, 137

Example 56

- A mixture of 2-acetylamino-5-(4-hydroxyphenylthio)thiazole (1.5 g) in a mixture of ethanol (40 ml) and aqueous 6N hydrochloric acid (6 ml) was refluxed for 4.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 10 using aqueous sodium hydroxide under ice cooling. The precipitates were collected by filtration and recrystallized from a mixture of ethanol and water (3:1) to give 2-amino-5-(4-hydroxyphenylthio)thiazole (1.05 g, yield : 84.0%).
- mp : 187-188 °C
 IR (Nujol) : 3450, 3350, 3200, 1625, 1600, 1500, 1320, 1245 cm⁻¹
 NMR (DMSO-d₆, 200MHZ, ppm) : 6.74 (2H, d, J=9Hz), 7.08 (2H, d, J=9Hz), 7.13 (1H, s), 7.34 (2H, s), 9.58 (1H, s)
- Mass : M⁺ 225, M 224, m/e 192, 182, 165, 137

Example 57

A mixture of 2-acetylamino-5-chlorothiazole (1.76 g), 4-methoxythiophenol (1.5 g) and potassium carbonate (2.0 g) in N,N-dimethylformamide (30 ml) was heated at 120 °C for 3.5 hours with stirring. The reaction mixture was poured into ice water and the precipitates were collected by filtration to give solid. The solid was recrystallized from ethanol to give 2-acetylamino-5-(4-methoxyphenylthio)thiazole (2.2 g, Yield : 78.6%).

mp : 190-191 °C

IR (Nujol) : 3175, 1695, 1565, 1490, 1295, 1250 cm⁻¹

¹H NMR (DMSO-d₆, 200MHZ, ppm) : 2.14 (3H, s), 3.74 (3H, s), 6.93 (2H, d, J=9Hz), 7.25 (2H, d, J=9Hz), 7.68 (1H, s), 12.31 (1H, s)

Mass : M⁺ 281, M 280, m/e 238, 205, 196, 151

Example 58

A mixture of 2-acetylamino-5-(4-methoxyphenylthio)thiazole (1.7 g) in a mixture of ethanol (40 ml) and aqueous 6N hydrochloric acid (6 ml) was refluxed for 4 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 10 using aqueous sodium hydroxide under cooling. The precipitates were collected by filtration, washed with water and recrystallized from ethanol to give 2-amino-5-(4-methoxyphenylthio)thiazole (1.25 g, yield : 86.8%).

mp : 119-120 °C

IR (Nujol) : 3400, 3275, 3100, 1635, 1595, 1520, 1460, 1240 cm⁻¹

¹H NMR (DMSO-d₆, 200MHZ, ppm) : 3.73 (3H, s), 6.91 (2H, d, J=9Hz), 7.17 (1H, s), 7.21 (2H, d, J=9Hz), 7.39 (2H, s)

Mass : M⁺ 239, M 238, m/e 206, 196, 151

Example 59

A mixture of 2-acetylamino-5-chlorothiazole (1.76 g), 5-mercapto-2-methyl-1,3,4-thiadiazole (1.3 g) and potassium carbonate (2.0 g) in N,N-dimethylformamide (40 ml) was heated at 120 °C for 4 hours with stirring. The reaction mixture was concentrated under reduced pressure and water was added to this residue. The mixture was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 150 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-acetylamino-5-(2-methyl-1,3,4-thiadiazol-5-ylthio)thiazole (1.65 g, yield : 60.7%).

mp : 242-244 °C

IR (Nujol) : 3250, 1695, 1550, 1300 cm⁻¹

¹H NMR (DMSO-d₆, 200MHZ, ppm) : 2.19 (3H, s), 2.63 (3H, s), 7.95 (1H, s), 12.58 (1H, s)

Mass : M⁺ 273, M 272, m/e 230, 188, 155, 131

Example 60

A mixture of 2-acetylamino-5-(2-methyl-1,3,4-thiadiazol-5-ylthio)thiazole (3.3 g) in a mixture of ethanol (70 ml), tetrahydrofuran (50 ml) and aqueous 6N-hydrochloric acid (200 ml) was refluxed for 6.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 8.5 using aqueous sodium bicarbonate and extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1). The organic layer was washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The organic solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 150 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-5-(2-methyl-1,3,4-thiadiazol-5-ylthio)thiazole (0.85 g, Yield : 58.2%).

mp : 203-205 °C (dec.)
 IR (Nujol) : 3450, 3300, 3100, 1640, 1520, 1485, 1220 cm⁻¹
 NMR (DMSO-d₆, 200MHZ, ppm) : 2.63 (3H, s), 7.42 (1H, s), 7.75 (2H, s)
 Mass : M⁺ 231, M 230, m/e 188, 154, 131, 113

5

Example 61

A mixture of 2-acetylamino-5-chlorothiazole (1.76 g), 5-mercapto-1-methyl-1H-tetrazole (1.2 g) and
 10 potassium carbonate (2 g) in N,N-dimethylformamide (40 ml) was heated at 130 °C for 3 hours with stirring.
 The reaction mixture was concentrated under reduced pressure and the residue was triturated with water.
 The precipitates were collected by filtration, washed with water and dried in vacuo to give solid. The solid
 was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 150g) and
 15 eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound
 were combined and concentrated under reduced pressure to give 2-acetylamino-5-(1-methyl-1H-tetrazol-5-
 ylthio)thiazole (2.1 g, yield : 82.0%).
 mp : 208-210 °C
 IR (Nujol) : 3450, 3250, 3150, 1690, 1665, 1550, 1295, 1225 cm⁻¹
 NMR (DMSO-d₆, 200MHZ, ppm) : 2.17 (3H, s), 4.11 (3H, s), 7.89 (1H, s), 12.51 (1H, s)
 20 Mass : M⁺ 257, M 256, m/e 214, 173, 159, 131

Example 62

A mixture of 2-acetylamino-5-(1-methyl-1H-tetrazol-5-ylthio)thiazole (2.0 g) in a mixture of ethanol (20
 25 ml) and aqueous 6N-hydrochloric acid (5 ml) was refluxed for 4 hours with stirring. The reaction mixture
 was concentrated under reduced pressure and the residue was adjusted to pH 8 using aqueous sodium
 bicarbonate under ice cooling. The precipitates were collected by filtration, washed with water and the solid
 was recrystallized from ethanol to give 2-amino-5-(1-methyl-1H-tetrazol-5-ylthio)thiazole (0.81 g, yield :
 30 48.5%).
 mp : 186-188 °C (dec.)
 IR (Nujol) : 3400, 3250, 3150, 1612, 1510, 1490, 1215 cm⁻¹
 NMR (DMSO-d₆, 200MHZ, ppm) : 4.03 (3H, s), 7.38 (1H, s), 7.63 (2H, s)
 Mass : M⁺ 215, M 214, m/e 131, 89, 83

35

Example 63

A mixture of 2-amino-5-bromothiazole hydrochloride (2.2 g), 4-amino-2-mercaptopyrimidine (2.2 g) and
 40 potassium carbonate (6.5 g) in N,N-dimethylformamide (100 ml) was heated at 90 °C for 2.5 hours with
 stirring. The reaction mixture was concentrated under reduced pressure and water was added to this
 residue. The solution was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with
 aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under
 reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel
 45 60, 70-230 mesh; Merck : 250 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions
 containing the objective compound were combined and concentrated under reduced pressure to give solid.
 The solid was triturated with ethanol to give 2-amino-5-(4-aminopyrimidin-2-ylthio)thiazole (1.25 g, yield :
 55.6%).
 mp : 185-187 °C (dec.)
 50 IR (Nujol) : 3450, 3300, 3175, 3100, 1645, 1630, 1580, 1545, 1340 cm⁻¹
 NMR (DMSO-d₆, 200MHZ, ppm) : 6.16 (1H, d, J=6Hz), 6.99 (2H, s), 7.07 (1H, s), 7.32 (2H, s), 7.86 (1H, d,
 J=6Hz)
 Mass : M⁺ 226, M 225, m/e 183, 139

55

Example 64

A mixture of 2-amino-5-bromo-4-methylthiazole hydrochloride (1.15 g), 2-mercaptopyrimidine (0.6 g)

- and potassium carbonate (1.7 g) in N,N-dimethylformamide (20 ml) was heated at 90 °C for 3.5 hours with stirring. The reaction mixture was poured into ice water. The mixture was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was
- 5 subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 100 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-4-methyl-5-(2-pyrimidinylthio)thiazole (0.65 g, yield : 58.0%).
- mp : 165-170 °C (dec.)
- 10 IR (Nujol) : 3300, 3175, 1630, 1555, 1490, 1320 cm⁻¹
- NMR (DMSO-d₆, 200MHZ, ppm) : 2.10 (3H, s), 7.24-7.33 (1H, m), 7.33 (2H, s), 8.64 (2H, d, J = 5Hz)
- Mass : M⁺ 225, M 224, m/e 209, 191, 182, 166, 145

15 Example 65

- A mixture of 2-amino-5-bromo-4-methylthiazole hydrochloride (4.5 g), 2-mercaptopyridine (2.3 g) and potassium carbonate (7.0 g) in N,N-dimethylformamide (100 ml) was heated at 90 °C for 3 hours with stirring. The reaction mixture was concentrated under reduced pressure and water was added to this
- 20 residue. The mixture was extracted with a mixture of tetrahydrofuran and ethyl acetate, washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 300 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give oil.
- 25 Again the oil was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 200 g) and eluted with a mixture of dichloromethane and acetone (5:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-4-methyl-5-(2-pyridylthio)thiazole (2.1 g, yield : 47.9%).
- NMR (DMSO-d₆, 200MHZ, ppm) : 2.13 (3H, s), 6.97 (1H, m), 7.15 (1H, m), 7.28 (2H, s), 7.65 (1H, m), 8.40
- 30 (1H, m)
- Mass : M⁺ 224, M 223, m/e 208, 190, 181, 145, 111

Example 66

- 35 A mixture of 2-amino-4-methyl-5-(2-pyridylthio)thiazole (1.7 g) and 3-chloroperbenzoic acid (1.8 g) in a mixture of chloroform (20 ml) and dichloromethane (50 ml) was stirred at 5 °C for 3.5 hours. The reaction mixture was washed with aqueous sodium bicarbonate and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on
- 40 silica gel (silica gel 60, 70-230 mesh; Merck : 100 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-4-methyl-5-(2-pyridylsulfinyl)thiazole (0.95 g, yield : 52.2 %).
- mp : 190-193 °C (dec.)
- NMR (DMSO-d₆, 200MHZ, ppm) : 2.38 (3H, s), 7.50-7.58 (1H, m), 7.70 (2H, s), 7.96 (1H, d, J = 8Hz), 8.07-
- 45 8.16 (1H, m), 8.6-8.63 (1H, m)
- Mass : M⁺ 240, M 239, m/e 223, 191, 161, 129, 111.

Example 67

- 50 A mixture of 2-acetylamino-5-bromothiazole (1.9 g), 2-mercaptoimidazole (0.9 g) and potassium carbonate (1.5 g) in N,N-dimethylformamide (30 ml) was heated at 90 °C for 2.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was extracted with methanol. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column
- 55 chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 150 g) and eluted with a mixture of chloroform and methanol (10:1). The fraction containing the objective compound were combined and concentrated under reduced pressure to give 2-acetylamino-5-(2-imidazolylthio)thiazole (1.8 g, yield : 87.4%).

mp : 230-235 °C (dec.)

IR (Nujol) : 3150, 3100, 1710, 1550, 1290 cm⁻¹

NMR (DMSO-d₆, 200MHZ, ppm) : 2.1 (3H, s), 6.80 (1H, s), 7.08 (2H, s), 7.6 (1H, s), 12.3 (1H, s) Mass : M⁺ 241, M 240, m/e 198, 156, 100

5

Example 68

A mixture of 2-acetylamino-5-(2-imidazolylthio)thiazole (1.8 g) in a mixture of concentrated hydrochloric acid (10 ml) and ethanol (50 ml) was refluxed for 5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 8.5 using sodium bicarbonate with cooling. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-(2-imidazolylthio)thiazole (0.35 g). The filtrate was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 100 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-5-(2-imidazolylthio)thiazole (0.55 g). Total amount of 2-amino-5-(2-imidazolylthio)thiazole was 0.90 g (yield : 60.4%).

mp : 209-211 °C (dec.)

IR (Nujol) : 3450, 3300, 1630, 1520, 1315 cm⁻¹ NMR: (DMSO-d₆, 200MHZ, ppm) : 7.05 (2H, s), 7.15 (1H, s), 7.36 (2H, s)

Mass : M⁺ 199, M 198, m/e 156, 139, 100

25

Example 69

A mixture of 2-acetylamino-5-bromothiazole (1.8 g), 3-hydroxy-2-mercaptopyridine (1.3 g) and potassium carbonate (2.0 g) in N,N-dimethylformamide (40 ml) was heated at 90 °C for 3.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The mixture was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 150 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-acetylamino-5-(3-hydroxypyridin-2-ylthio)thiazole (2.4 g, yield : 89.9%).

mp : 236-238 °C (dec.)

IR (Nujol) : 3175, 1690, 1565, 1300 cm⁻¹

NMR (DMSO-d₆, 200MHZ, ppm) : 2.16 (3H, s), 7.02-7.17 (3H, m), 7.58 (1H, s), 7.83 (1H, d, J=6Hz), 10.70 (1H, s), 12.30 (1H, s)

Mass : M⁺ 268, M 267, m/e 225, 183, 127

Example 70

A mixture of 2-acetylamino-5-(3-hydroxypyridin-2-ylthio)thiazole (2 g) in a mixture of ethanol (40 ml), tetrahydrofuran (20 ml) and aqueous 6N-hydrochloric acid (13 ml) was refluxed for 5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 8.5 using aqueous sodium bicarbonate and extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1). The organic layer was washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give 2-amino-5-(3-hydroxypyridin-2-ylthio)thiazole (1.15 g, yield : 68.9%).

mp : 128-130 °C

IR (Nujol) : 3500, 3400, 3300, 1640, 1570, 1520, 1500, 1330, 1200 cm⁻¹

NMR (DMSO-d₆, 200MHZ, ppm) : 6.97-7.11 (3H, m), 7.29 (2H, s), 7.82 (1H, d, J=6Hz), 10.57 (1H, s)

Mass : M⁺ 226, M 225, m/e 183, 139, 100

Example 71

To a mixture of 2-amino-5-(3-hydroxypyridin-2-ylthio)thiazole (4.6 g) in a mixture of chloroform (100 ml), dichloromethane (200 ml) and N,N-dimethylformamide (50 ml) was added dropwise the solution of 3-chloroperbenzoic acid (4.3 g) in chloroform (50 ml) at 5 °C with stirring. The mixture was stirred at room temperature for 5 hours. The reaction mixture was extracted with aqueous diluted hydrochloric acid and the aqueous layer was washed with ethyl acetate. The aqueous layer was adjusted to pH 5.7 using sodium bicarbonate and extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1). The organic layer was washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 250 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-5-(3-hydroxypyridin-2-ylsulfanyl)thiazole (0.65 g, yield : 12.2%).

mp : 155-158 °C (dec.)

IR (Nujol) : 3300, 3150, 1620, 1565, 1515, 1300 cm⁻¹

NMR (DMSO-d₆, 200MHZ, ppm) : 7.29-7.40 (2H, m), 7.59 (1H, s), 7.75 (2H, s), 8.15 (1H, br s)

Mass : m/e 225, 220, 205

Example 72

A mixture of 2-acetylamino-5-bromothiazole (1 g), 3-mercaptopyridine hydrochloride (1 g) and potassium carbonate (1.5 g) in N,N-dimethylformamide (10 ml) was heated at 90 °C for 4.5 hours with stirring. The reaction mixture was poured into ice water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-acetylamino-5-(3-pyridylthio)thiazole (0.9 g, yield : 81.8%).

mp : 203-205 °C (dec.)

IR (Nujol) : 3170, 1700, 1570, 1300 cm⁻¹

NMR (DMSO-d₆, 200MHZ, ppm) : 2.16 (3H, s), 7.33-7.39 (1H, s), 7.60 (1H, d, J=8Hz), 7.81 (1H, s), 8.42-8.54 (2H, m), 12.45 (1H, s)

Mass : M⁺ 252, M 251, m/e 209, 176, 167, 111

Example 73

A mixture of 2-acetylamino-5-(3-pyridylthio)thiazole (8.5 g) in a mixture of ethanol (160 ml), tetrahydrofuran (50 ml) and aqueous 6N hydrochloric acid (100 ml) was refluxed for 4 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 8.5 using aqueous sodium bicarbonate and the precipitates were collected by filtration, washed with water and dried in vacuo to give 2-amino-5-(3-pyridylthio)thiazole (5.6 g yield : 78.9%).

mp : 140-142 °C

IR (Nujol) : 3400, 3300, 3125, 1630, 1530, 1490 cm⁻¹

NMR (DMSO-d₆, 200MHZ, ppm) : 7.30 (1H, s), 7.32-7.39 (1H, m), 7.55 (2H, s), 7.57-7.61 (1H, m), 8.40 (2H, d, J=7Hz)

Mass : M⁺ 210, M 209, m/e 167, 122, 99

Example 74

To a mixture of 2-amino-5-(3-pyridylthio)thiazole (3.0 g) in a mixture of dichloromethane (100 ml) and chloroform (100 ml) was added dropwise the solution of 3-chloroperbenzoic acid (3.4 g) in dichloromethane (50 ml) at 5 °C with stirring. The mixture was stirred at 5 °C for 3 hours. The reaction mixture was washed with aqueous sodium bicarbonate and the aqueous layer was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with aqueous saturated sodium chloride and dried over magnesium sulfate. The solvent was concentrated under reduced pressure and the residue was recrystallized from ethanol to give 2-amino-5-(3-pyridylsulfanyl)thiazole (1.2 g, yield : 37.2%).

mp : 178-179 °C

IR (Nujol) : 3300, 3150, 1630, 1580, 1520, 1485, 1325, 1220 cm⁻¹

NMR (DMSO- d_6 , 200MHZ, ppm) : 7.58-7.70 (1H, m), 7.80 (1H, s), 8.00 (2H, s), 7.98-8.05 (1H, m), 8.72 (2H, br s)

Mass : M^+ 226, M 225, m/e 209, 177, 147

5

Example 75

A mixture of 2-amino-5-(3-pyridylsulfinyl)thiazole (1.6 g) and 3-chloroperbenzoic acid (1.8 g) in a mixture of chloroform (150 ml), dichloromethane (50 ml) and N,N-dimethylformamide (5 ml) was stirred at room temperature for 3 hours. The reaction mixture was extracted with diluted hydrochloric acid and washed with ethyl acetate. The aqueous layer was adjusted to pH 8.5 using sodium bicarbonate and the mixture was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1). The organic layer was washed with water and dried over magnesium sulfate. The solvent was concentrated under reduced pressure and the residue was triturated with ethanol to give 2-amino-5-(3-pyridylsulfonyl)thiazole (0.30 g, yield : 17.5%).

mp : 218-220 °C (dec.)

IR (Nujol) : 3420, 3300, 1650, 1520, 1310 cm^{-1}

NMR (DMSO- d_6 , 200MHZ, ppm) : 7.63-7.70 (1H, m), 7.78 (1H, s), 8.20 (2H, s), 8.27 (1H, d, J=8Hz), 8.85 (1H, d, J=4Hz), 9.06 (1H, s)

Mass : M^+ 242, M 241, m/e 177, 135, 99

Example 76

A mixture of 2-acetylamino-5-bromothiazole (2.2 g), 2-mercapto-5-trifluoromethylpyridine (1.9 g) and potassium carbonate (2.0 g) in N,N-dimethylformamide (40 ml) was heated at 90 °C for 4.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The precipitates were collected by filtration, washed with water and dried in vacuo to give 2-acetylamino-5-(5-trifluoromethylpyridin-2-ylthio)thiazole (3.2 g, yield : 100%).

mp : 165-170 °C (dec.)

IR (Nujol) : 3175, 1695, 1640, 1600, 1565, 1330 cm^{-1}

NMR (DMSO- d_6 , 200MHZ, ppm) : 2.19 (3H, s), 7.21 (1H, d, J=12Hz), 7.83 (1H, s), 8.04-8.15 (1H, m), 8.18 (1H, br s), 12.53 (1H, s)

Mass : M^+ 321, M 320, M319, m/e 277, 235, 191

35

Example 77

A mixture of 2-acetylamino-5-(5-trifluoromethylpyridin-2-ylthio)thiazole (3.2 g) in a mixture of ethanol (60 ml), tetrahydrofuran (30 ml) and aqueous 6N-hydrochloric acid (10 ml) was refluxed for 3 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in water. The solution was adjusted to pH 8.5 using aqueous sodium bicarbonate and extracted with mixture of tetrahydrofuran and ethyl acetate (1:1) and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 150 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-5-(5-trifluoromethylpyridin-2-ylthio)thiazole (2.1 g, yield : 75.8%).

mp : 135-138 °C

IR (Nujol) : 3400, 3300, 3100, 1640, 1600, 1560, 1520, 1330 cm^{-1}

NMR (DMSO- d_6 , 200MHZ, ppm) : 7.26 (1H, d, J=8Hz), 7.33 (1H, s), 7.66 (2H, s), 7.81 (1H, d, J=8Hz), 8.80 (1H, s)

Mass : M^+ 278, M 277, m/e 235, 191, 146, 131

Example 78

To a solution of 2-amino-5-(5-trifluoromethylpyridin-2-ylthio)thiazole (0.6 g) in dichloromethane (20 ml) was added portionwise 3-chloroperbenzoic acid (0.6 g) at 5 °C with stirring. The mixture was stirred at 5 °C

for 3 hours. The reaction mixture was washed with aqueous sodium bicarbonate and dried over magnesium sulfate. The solvent was concentrated under reduced pressure to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 30 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-5-(5-trifluoromethylpyridin-2-ylsulfanyl)thiazole (0.52 g, yield : 81.9%).
 mp : 144-145 °C
 NMR (DMSO- d_6 , 200MHZ, ppm) : 7.82 (1H, s), 7.92 (2H, s), 8.21 (1H, d, J = 8Hz), 8.56 (1H, d, J = 8Hz), 9.08 (1H, s)
 10 Mass : M 293, m/e 277, 245, 226, 179, 147

Example 79

15 A mixture of 2-acetylamino-5-bromothiazole (2.2 g), 4-amino-2-mercaptopyrimidine (1.3 g) and potassium carbonate (2.0 g) in N,N-dimethylformamide (50 ml) was heated at 90 °C for 2 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was triturated with water. The precipitation was collected by filtration, washed with water and dried in vacuo to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 200 g) and eluted
 20 with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-acetylamino-5-(4-aminopyrimidin-2-ylthio)thiazole (1.3 g, yield : 48.7%).
 mp : 255-258 °C (dec.)
 IR (Nujol) : 3400, 3350, 3200, 1692, 1650, 1585, 1325, 1300 cm^{-1}
 25 NMR (DMSO- d_6 , 200MHZ, ppm) : 2.16 (3H, s), 6.17 (1H, d, J = 6Hz), 7.02 (2H, s), 7.59 (1H, s), 7.85 (1H, d, J = 6Hz), 12.31 (1H, s)
 Mass : M⁺ 268, M 267, m/e 225, 205, 183

30 Example 80

Starting from 2-acetylamino-5-bromothiazole, 2-acetylamino-5-(4-hydroxypyrimidin-2-ylthio)thiazole (0.35 g, yield : 28.8%) was obtained according to a similar manner to that of Example 67.
 IR (Nujol) : 3150, 1665, 1565, 1535, 1300, 1275, 1230 cm^{-1}
 35 NMR (DMSO- d_6 , 200MHZ, ppm) : 2.18 (3H, s), 6.25 (1H, d, J = 6Hz), 7.70 (1H, s), 7.93 (1H, d, J = 6Hz), 12.0-12.6 (2H, m)
 Mass : M⁺ 269, M 268, m/e 259, 197, 135

40 Example 81

A mixture of 2-acetylamino-5-(4-hydroxypyrimidin-2-ylthio)thiazole (3.7 g) in a mixture of ethanol (100 ml), tetrahydrofuran (40 ml) and aqueous 6N-hydrochloric acid (20 ml) was refluxed for 6.5 hours with stirring. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in
 45 water. The solution was adjusted to pH 8.5 using aqueous sodium bicarbonate. The precipitates were collected by filtration, washed with water and dried in vacuo to give solid. The solid was subjected to column chromatography on silica gel (silica gel 60, 70-230 mesh; Merck : 250 g) and eluted with a mixture of chloroform and methanol (10:1). The fractions containing the objective compound were combined and concentrated under reduced pressure to give 2-amino-5-(4-hydroxypyrimidin-2-ylthio)thiazole (0.45 g, yield :
 50 14.5%).
 mp : 210-220 °C (dec.)
 IR (Nujol) : 3450, 3350, 3125, 1675, 1510, 1230 cm^{-1}
 NMR (DMSO- d_6 , 200MHZ, ppm) : 5.45 (1H, d, J = 7Hz), 7.30-7.40 (3H, m), 7.64 (1H, m)
 Mass : m/e 220, 205, 132, 112
 55

Example 82

Starting from 2-acetylamino-5-bromothiazole, 2-acetylamino-5-(4-methylpyrimidin-2-ylthio)thiazole (2.89 g, yield : 48.0%) was obtained according to a similar manner to that of Example 67.

mp : 210 °C (dec.)

IR (Nujol) : 3170, 1720, 1695, 1575, 1555, 1335 cm⁻¹

5 NMR (DMSO-d₆, 200MHZ, ppm) : 2.19 (3H, s), 2.39 (3H, s), 7.16 (1H, d, J=5Hz), 7.70 (1H, s), 8.46 (1H, d, J=5Hz), 12.38 (1H, s)

Mass : M⁺ 268, M⁺ 267, M 266, m/e 224, 182, 165

10 Example 83

Starting from 2-acetylamino-5-(4-methylpyrimidin-2-ylthio)thiazole, 2-amino-5-(4-methylpyrimidin-2-ylthio)thiazole (0.40 g, yield : 16.4%) was obtained according to a similar manner to that of Example 58.

mp : 158-159 °C

15 IR (Nujol) : 3430, 3280, 3100, 1620, 1565, 1520, 1490, 1330, 1210 cm⁻¹

NMR (DMSO-d₆, 200MHZ, ppm) : 2.39 (3H, s), 7.14 (1H, d, J=5Hz), 7.15 (1H, s), 7.43 (2H, s), 8.47 (1H, d, J=5Hz)

Mass : M⁺ 226, M⁺ 225, M 224, m/e 182, 138

20

| Analysis Calcd. for C ₈ H ₈ N ₄ S ₂ | |
|---|--|
| Found : | C 42.84, H 3.59, N 24.99 C 42.81, H 3.50, N 24.86 |

25

Example 84

30 To a solution of 2-amino-5-(2-pyrimidinylthio)thiazole (1.0 g) in a pyridine (20 ml) was dropwise added methanesulfonyl chloride (0.8 ml) at 5 °C with stirring. The mixture was stirred at room temperature for 24 hours. The reaction mixture was concentrated under reduced pressure and water was added to this residue. The mixture was extracted with a mixture of tetrahydrofuran and ethyl acetate (1:1), washed with aqueous
35 saturated sodium chloride and dried over magnesium sulfate. The solution was concentrated under reduced pressure to give solid. The solid was recrystallized from 50% ethanol to give 2-methanesulfonylamino-5-(2-pyrimidinylthio)thiazole (0.60 g, yield : 43.8%).

mp : 200 °C (dec.)

IR (Nujol) : 3120, 1585, 1545, 1440, 1305, 1140 cm⁻¹

40 NMR (DMSO-d₆, 200MHZ, ppm) : 2.98 (3H, s), 7.35 (1H, t, d=7Hz), 7.76 (1H, s), 8.71 (2H, d, J=7Hz), 12.89 (1H, br s)

Mass : M⁺ 290, M⁺ 289, M 288, m/e 209, 168

45

| Analysis Calcd. for C ₈ H ₈ N ₄ O ₂ S ₃ | |
|--|--|
| Found : | C 33.32, H 2.80, N 19.43 C 33.04, H 2.74, N 19.06 |

50

Example 85

Starting from 2-amino-5-(2-pyrimidinylthio)thiazole, 2-amino-5-(2-pyrimidinylsulfinyl)thiazole (0.52 g, yield : 32.2%) was obtained according to a similar manner to that of Example 66.

mp : 206 °C (dec.)

IR (Nujol) : 3300, 3200, 1615, 1565, 1545, 1520, 1230, 1150 cm⁻¹

NMR (DMSO-d₆ 200MHZ ppm) : 7.67 (1H, t J=5Hz) 7.73 (1H,s), 7.87 (2H, s), 8.99 (2H, d, J=5Hz)

Mass : M 226, m/e 210, 178, 168, 147

Analysis Calcd. for $C_7H_6N_4OS_2$

| | |
|---------|--------------------------|
| | C 37.16, H 2.67, N 24.76 |
| Found : | C 36.78, H 2.62, N 24.62 |

Example 86


Starting from 2-amino-5-(2-pyrimidinylthio)thiazole, 2-amino-5-(2-pyrimidinylsulfonyl)thiazole (0.464 g, yield : 8.1%) was obtained according to a similar manner to that of Example 75.

mp : 214 °C (dec.)


IR (Nujol) : 3400, 3100, 1615, 1570, 1515, 1335, 1210, 1140 cm^{-1} NMR (DMSO- d_6 , 200MHZ, ppm) : 7.73 (1H, s), 7.80 (1H, t, d = 5Hz), 8.23 (2H, s), 9.05 (2H, d, J = 5Hz)Mass : M^{+2} 244, M^{+1} 243, M 242, m/e 178, 136

Claims

1. A compound of the formula :



wherein

 R^1 is hydrogen or acyl which may be substituted with halogen, R^2 is hydrogen, lower alkyl, hydroxy(lower)alkyl, halogen or carboxy,A is $-CH_2-$, $-CO-$, $-C(=NOR^4)-$ [wherein R^4 is lower alkyl],

[wherein m is 0, 1 or 2], and

 R^3 is aryl which may be substituted with halogen, hydroxy, lower alkoxy, nitro, amino or acylamino; or

N-containing unsaturated heterocyclic group which may be substituted with lower alkyl, amino, hydroxy or halo(lower)alkyl,

and pharmaceutically acceptable salts thereof.

2. A compound of claim 1, wherein R^1 is hydrogen, lower alkanoyl, lower alkanesulfonyl or benzoyl which may be substituted with halogen,3. A compound of claim 2, wherein R^3 is phenyl which may be substituted with halogen, hydroxy, lower alkoxy, nitro, amino, lower alkanoylamino or lower alkanesulfonylamino; or pyridyl, pyrimidinyl, imidazolyl, tetrazolyl or thiadiazolyl, each of which may be substituted with lower alkyl, amino, hydroxy or halo(lower)alkyl.

4. A compound of claim 3, wherein

 R^1 is hydrogen.

5. A compound of claim 4, wherein

 R^2 is hydrogen.

6. A compound of claim 5, wherein

A is




5 [wherein m is 0, 1 or 2].

6. A compound of claim 5, wherein

R³ is pyridyl, pyrimidinyl, imidazolyl, tetrazolyl or thiadiazolyl.

7. A process for preparing a compound of the formula :

10



15


wherein

R¹ is hydrogen or acyl which may be substituted with halogen,

20 R² is hydrogen, lower alkyl, hydroxy(lower)alkyl, halogen or carboxy,

A is -CH₂-, -CO-, -C(=NOR⁴)- [wherein R⁴ is lower alkyl],

25




[wherein m is 0, 1 or 2], and

30 R³ is aryl which may be substituted with halogen, hydroxy, lower alkoxy, nitro, amino or acylamino; or N-containing unsaturated heterocyclic group which may be substituted with lower alkyl, amino, hydroxy or halo(lower)alkyl,

or its salt, which comprises

(1) deacylating a compound of the formula :


35



40

or its salt to give a compound of the formula :

45



50


or its salt, in which R², R³ and A are each as defined above,

55

R² is acyl which may be substituted with halogen, or

(2) reacting a compound of the formula :


5



10

or its salt with a compound of the formula :
or its salt to give a compound of the formula :

15




20

or its salt, in which R^1 , R^2 and R^3 are each as defined above,
 X is halogen and l is 0 or 1, or
(3) reducing a compound of the formula :

25


30



35

or its salt to give a compound of the formula :


40



45


or its salt, in which R^1 , R^2 and A are each as defined above,
or
(4) subjecting a compound of the formula :

50




55


or its salt to oxidation to give a compound of the formula :




15 or its salt, in which R^1 , R^2 , R^3 and l are each as defined above, n is 0 or 1 and q is 1 or 2, provided that q is 2 when n is 1, or
(5) subjecting a compound of the formula :




25 or its salt to oxidation to give a compound of the formula :




35 or its salt, in which R^1 and R^3 are each as defined above,
or
(6) halogenating a compound of the formula :




or its salt to give a compound of the formula :




55 or its salt, in which R^1 , R^3 , A and X are each as defined above, or
(7) acylating a compound of the formula :




10 or its reactive derivative at the amino group or a salt thereof to give a compound of the formula :



20 or its salt, in which R², R³, R⁵ and A are each as defined above, or
(8) acylating a compound of the formula :



or its reactive derivative at the amino group or a salt thereof to give a compound of the formula :




or its salt, in which R¹, R² and A are each as defined above,
R⁶ is acylamino, or


(9) reacting a compound of the formula :

45 R³-CH₂-CO-CH₃

or its salt with a compound of the formula :



or its salt to give a compound of the formula :




or its salt, in which R¹ and R³ are each as defined above,

or


(10) reacting a compound of the formula :



or its salt with a compound of the formula :




or its salt to give a compound of the formula :



or its salt, in which R¹, R³ and X are each as defined above,

or


(11) reacting a compound of the formula :



or its salt with a compound of the formula :



or its salt to give a compound of the formula :



or its salt, in which R¹, R², R³ and R⁴ are each as defined above.

55 8. A pharmaceutical composition which comprises, as an active ingredient, a compound of claim 1 or a pharmaceutically acceptable salt thereof in admixture with pharmaceutically acceptable carriers.

9. Use of a compound of claim 1 or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for therapeutic treatment of rheumatism, nephritis, thrombocytopenia, tumor and side effects of

an antitumor agent.

10. A compound of claim 1 or a pharmaceutically acceptable salt thereof for use as a medicament.

5

10

15

20

25

30

35

40

45

50

55



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 412 404 A3**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90114628.2

(51) Int. Cl.⁵: **C07D 277/38, A61K 31/425,
C07D 277/56, C07D 417/12**

(22) Date of filing: 31.07.90

(30) Priority: 07.08.89 GB 8918045
21.02.90 GB 9003930

(43) Date of publication of application:
13.02.91 Bulletin 91/07

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of the search report:
12.06.91 Bulletin 91/24

(71) Applicant: **FUJISAWA PHARMACEUTICAL CO.,
LTD.**
4-7, Doshomachi 3-chome Chuo-ku
Osaka-shi Osaka 541(JP)


(72) Inventor: Matsuo, Masaaki

**4-12, Nakasakurazuka 5-chome
Toyonaka-shi, Osaka 560(JP)**
Inventor: **Ogino, Takashi**
**4-6-208, Minamigoyo 2-chome
Kita-ku, Kobe-shi, Hyogo 651-11(JP)**
Inventor: **Igari, Norihiro**
**2-12-401 Ohsho-kita 4-chome
Amagasaki-shi, Hyogo 660(JP)**
Inventor: **Seno, Hachiro**
**12-1, Sengokuhigashimachi
Kadoma-shi, Osaka 571(JP)**
Inventor: **Shimomura, Kyoichi**
**3-9, Tsugumodai 2-chome
Suita-shi, Osaka 565(JP)**

(74) Representative: **Türk, Gille, Hrabal**
Brucknerstrasse 20
W-4000 Düsseldorf 13(DE)

(54) **Thiazole derivatives, processes for production thereof and pharmaceutical compositions comprising the same.**

(57) A compound of the formula :




wherein

R¹ is hydrogen or acyl which may be substituted with halogen,

R² is hydrogen, lower alkyl, hydroxy(lower)alkyl, halogen or carboxy,

A is -CH₂-, -CO-, -C(=NOR⁴)- [wherein R⁴ is lower alkyl,



[wherein m is 0, 1 or 2], and

R³ is aryl which may be substituted with halogen, hydroxy, lower alkoxy, nitro, amino or acylamino; or N-containing unsaturated heterocyclic group which may be substituted with lower alkyl, amino, hydroxy or halo(lower)alkyl,

and pharmaceutically acceptable salts thereof, processes for their preparation and pharmaceutical compositions comprising them as an active ingredient in admixture with pharmaceutically acceptable carriers.

EP 0 412 404 A3



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 11 4628

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|---|--|--|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int. Cl.5) |
| X | JOURNAL OF THE INDIAN CHEMICAL SOCIETY, vol. 39, no. 6, 1962, pages 420-426, Calcutta, IN; C.S. MAHAJAN-SHETTI et al.: "2-Aminothiazole derivatives. Part I" * The whole article * | 1-5,7-10 | C 07 D 277/38 A 61 K 31/425 C 07 D 277/56 C 07 D 417/12 |
| X | JOURNAL OF THE INDIAN CHEMICAL SOCIETY, vol. 40, no. 11, 1963, pages 921-924, Calcutta, IN; C.S. MAHAJAN-SHETTI et al.: "2-Aminothiazole derivatives. Part III" * The whole article * | 1-5,7-10 | |
| X | CHEMICAL ABSTRACTS, vol. 82, no. 13, 31st March 1975, page 93, abstract no. 81110d, Columbus, Ohio, US; C.S. MAHAJANSHETTI et al.: "2-Amino-4-methyl-5-thiazolyl phenyl sulfides and sulfones as possible antibacterials", & J. KARNATAK UNIV. 1973, 18, 5-10 * Abstract * | 1-3,5,7-10 | |
| X | JOURNAL OF MEDICINAL CHEMISTRY, vol. 12, no. 5, May 1969, pages 357-363, Washington, DC, US; E.F. ELSLAGER et al.: "Repository drugs. IV. 4',4'''-sulfonylbisacetanilide (acedapsona, DADDS) and related sulfanilylanilides with prolonged antimalarial and antileprotic action" * Page 359, compound X * | 1,2 | |
| X | CANADIAN JOURNAL OF CHEMISTRY, vol. 45, no. 15, 1967, pages 1807-1810, CA; C.S. MAHAJANSHETTI et al.: "A facile nucleophilic displacement of bromine in 2-amino-4-methyl-5-bromothiazole by a thiophenoxide anion" * The whole article * | 1,2 | |
| X | ARZNEIMITTEL-FORSCHUNG, vol. 26, no. 8, August 1976, pages 1343-1547, Aulendorf, DE; I. HAGEDORN et al.: "Reaktivierung phosphorylierter Acetylcholinesterase (AChE): Quartäre Salze vinyloger Pyridinaldoxime" * The whole article * | 1,2,7-10 | |
| The present search report has been drawn up for all claims | | | |
| Place of search | | Date of completion of search | Examiner |
| The Hague | | 08 March 91 | HENRY J.C. |
| CATEGORY OF CITED DOCUMENTS | | | |
| X: particularly relevant if taken alone | | E: earlier patent document, but published on, or after the filing date | |
| Y: particularly relevant if combined with another document of the same category | | D: document cited in the application | |
| A: technological background | | L: document cited for other reasons | |
| O: non-written disclosure | | | |
| P: intermediate document | | 8: member of the same patent family, corresponding document | |
| T: theory or principle underlying the invention | | | |



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 11 4628

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|---|---|---|---|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int. Cl.5) |
| X | FR-A-1 321 897 (SANKYO) * Abstract; example 3a, page 2 * | 1 | |
| X | EP-A-0 117 082 (FUJISAWA) * Claims * | 1,7-10 | |
| D,X | MONATSHEFTE FÜR CHEMIE, vol. 120, no. 11, November 1989, pages 991-995, Springer-Verlag, AT; M.M. EL-KERDAWY et al.: "Synthesis of substituted 4H-thiazolo[4,5-b][1]benzothiopyran-4-ones as possible schistosomicidal agents" * Page 992, compounds 4,5,7, page 994 * | 1,2 | |
| | | | TECHNICAL FIELDS SEARCHED (Int. Cl.5) |
| The present search report has been drawn up for all claims | | | |
| Place of search | | Date of completion of search | Examiner |
| The Hague | | 08 March 91 | HENRY J.C. |
| <p>CATEGORY F CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone</p> <p>Y: particularly relevant if combined with another document of the same category</p> <p>A: technological background</p> <p>O: non-written disclosure</p> <p>P: intermediate document</p> <p>T: theory or principle underlying the invention</p> | | <p>E: earlier patent document, but published on, or after the filing date</p> <p>D: document cited in the application</p> <p>L: document cited for other reasons</p> <p>&: member of the same patent family, corresponding document</p> | |